




**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

REC'D 24 JUL 2003

Patent Office PCT
Canberra

I, JULIE BILLINGSLEY, TEAM LEADER EXAMINATION SUPPORT AND
SALES hereby certify that annexed is a true copy of the Provisional specification
in connection with Application No. PS 3339 for a patent by THE
AUSTRALIAN NATIONAL UNIVERSITY as filed on 02 July 2002.



WITNESS my hand this
Fourteenth day of July 2003

A handwritten signature in cursive script, appearing to read 'J. Billingsley'.

JULIE BILLINGSLEY
TEAM LEADER EXAMINATION
SUPPORT AND SALES

AUSTRALIA

Patents Act 1990

The Australian National University

PROVISIONAL SPECIFICATION

Invention Title:

Method of producing plants having enhanced transpiration efficiency and plants produced therefrom I

The invention is described in the following statement:

FIELD OF THE INVENTION

The present invention relates to the field of plant breeding and the production of genetically engineered plants. More specifically, the invention described herein provides genes that are capable of enhancing the transpiration efficiency of a plant
5 when expressed therein. These genes are particularly useful for the production of plants having enhanced transpiration efficiency, by both traditional plant breeding and genetic engineering approaches. The invention further extends to plants produced by the methods described herein.

10 BACKGROUND TO THE INVENTION

1. General

This specification contains nucleotide and amino acid sequence information prepared using PatentIn Version 3.1, presented herein after the claims. Each nucleotide sequence is identified in the sequence listing by the numeric indicator <210> followed by the sequence identifier (e.g. <210>1, <210>2, etc). The length
15 and type of sequence (DNA, protein (PRT), etc), and source organism for each nucleotide sequence, are indicated by information provided in the numeric indicator fields <211>, <212> and <213>, respectively. Nucleotide sequences referred to in the specification are defined by the term "SEQ ID NO:", followed by the sequence identifier (eg. SEQ ID NO: 1 refers to the sequence in the sequence listing designated
20 as <400>1).

The designation of nucleotide residues referred to herein are those recommended by the IUPAC-IUB Biochemical Nomenclature Commission, wherein A represents Adenine, C represents Cytosine, G represents Guanine, T represents thymine, Y
25 represents a pyrimidine residue, R represents a purine residue, M represents Adenine or Cytosine, K represents Guanine or Thymine, S represents Guanine or Cytosine, W represents Adenine or Thymine, H represents a nucleotide other than Guanine, B represents a nucleotide other than Adenine, V represents a nucleotide other than Thymine, D represents a nucleotide other than Cytosine and N represents
30 any nucleotide residue.

As used herein the term "derived from" shall be taken to indicate that a specified integer is obtained from a particular source albeit not necessarily directly from that source.

5 Throughout this specification, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated step or element or integer or group of steps or elements or integers but not the exclusion of any other step or element or integer or group of elements or integers.

10

Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood that the invention includes all such variations and modifications. The invention also includes all of the steps, features, compositions and compounds
15 referred to or indicated in this specification, individually or collectively, and any and all combinations or any two or more of said steps or features.

The present invention is not to be limited in scope by the specific embodiments described herein, which are intended for the purposes of exemplification only.

20 Functionally equivalent products, compositions and methods are clearly within the scope of the invention, as described herein.

2. Description of the related art

It is well known that virtually all plants require a certain quantity of water for proper growth and development, because CO₂ fixation and photosynthate assimilation by plants cost water. A significant quantity of water absorbed by plants from the soil
25 returns to the atmosphere *via* plant transpiration.

Transpiration efficiency is a measure of the amount of dry matter produced by a plant per unit of water transpired, or, in other words, carbon gain relative to water lost through transpiration.

30

For plants having low transpiration efficiency, or when water is in short supply, the loss of water through transpiration can limit key metabolic processes associated with

plant growth and development. For example, during drought, or when plants having low transpiration efficiency are grown in arid and semi-arid environments, plant productivity as determined by dry matter production or photosynthetic rate, is considerably reduced. Accordingly, the production of plants having enhanced water use efficiency or transpiration efficiency is highly desirable for their adaptation to arid or semi-arid conditions, or to enhance their drought resistance.

The enhancement of water use efficiency or transpiration efficiency by plants is also highly desirable in consideration of global climatic change and increasing pressure on world water resources. The inefficient utilization of agricultural water is known to impact adversely upon the supply of navigable water, potable water, and water for industrial or recreational use. Accordingly, the production of plants having enhanced transpiration efficiency is highly desirable for reducing the pressure on these water resources. It is also desirable for increasing plant productivity under well-watered conditions.

By enhancing transpiration efficiency, carbon gain rates are enhanced per unit of water transpired, thereby stimulating plant growth under well-watered conditions, or alternatively, under mild or severe drought conditions. This is achieved by enhancing carbon gain more than transpiration rate, or by reducing the amount of water lost at any particular rate of carbon fixation. Those skilled in the art also consider that for a given growth rate plants having enhanced transpiration efficiency dry out soils more slowly, and use less water, than less efficient near-isogenic plants.

Several chemical as well as environmental pre-treatments have been described for enhancing the ability of plant seedlings to survive drought, either by reducing transpiration or by reducing the amount of water that is actually lost to the atmosphere.

Known environmental treatments largely involve the use of physical barriers. Whilst placing a physical barrier over plant stomata is known to reduce water loss via transpiration, the procedure is not always desirable or practicable for field-grown crops. For example, physical barriers over plant stomata may inhibit certain gas-exchange processes of the plant. It is more desirable to enhance actual transpiration

efficiency or water use efficiency of the plant through manipulation of intrinsic plant function.

Chemical agents are typically the so-called "anti-transpirant" or "anti-desiccant" agents, both of which are applied to the leaves. Anti-transpirants are typically films or metabolic anti-transpirants.

These products form a film on leaves, thereby either blocking stomatal pores, or coating leaf epidermal cells with a water-proof film. Typical film anti-transpirants include waxes, wax-oil emulsions, higher alcohols, silicones, plastics, latexes and resins. For example, Elmore, United States Patent No. 4,645,682 disclosed an anti-transpirant consisting of an aqueous paste wax; Cushman *et al.*, United States Patent Nos. 3,791,839 and 3,847,641 also disclosed wax emulsions for controlling transpiration in plants; and Petrucco *et al.*, United States Patent No. 3,826,671, disclosed a polymer composition said to be effective for controlling transpiration in plants.

Metabolic anti-transpirants generally close stomata, thereby reducing the rate of transpiration. Typical metabolic anti-transpirants include succinic acids, phenylmercuric acetate, hydroxysulfonates, the herbicide atrazine, sodium azide, and phenylhydrazones, as well as carbon cyanide.

Compounds having plant growth regulator activity have also been shown to be useful for reducing transpiration. For example, Bliesner *et al.*, United States Patent No. 4,671,816, disclosed an acetylene compound, said to possess utility for regulating plant growth, whilst Kuznetsov *et al.* (Russian Patent No. SU 1,282,492; .. Russian Patent Application No. SU 1,253,559-A1), and Smirnov *et al* (Russian Patent No. SU 1,098,934) disclosed the use of derivatives of 2-methyl-5-hydroxybenzimidazole, and the chloride or bromide salts thereof, as anti-transpirant growth regulators. Vichnevetskaia (USSN 5,589,437 issued December 31, 1996) also describe hydroxybenzimidazole derivatives for enhancing the drought resistance of plants by reducing transpiration, however have the advantage of being applicable to plant seed or roots. Schulz *et al.*, United States Patent No. 4,943,315, also disclosed formulations comprising an acetylene and a phenylbenzylurea compound, for

reducing transpiration in plants and/or for avoiding impairment to plants caused by heat and dry conditions. Abscissic acid has also been shown to reduce or suppress transpiration in plants (eg. *Helv. Chim. Acta*, 71, 931, 1988; *J. Org. Chem.*, 54, 681, 1989; and Japanese Patent Publication No. 184,966/1991).

5

Metabolic anti-transpirants are costly to produce and often exhibit phytotoxic effects or inhibit plant growth Kozlowski (1979), *In: Tree Growth and Environmental Stresses* (Univ. of Washington Press, Seattle and London), and are not practically used.

10

Recent studies have examined alternative methods for enhancing transpiration efficiency, particularly breeding approaches to select lines that grow more efficiently under mild drought conditions. Carbon isotope discrimination has been used to identify *Arabidopsis* ecotypes with contrasted transpiration efficiencies (Masle *et al.*, *In: Stable isotopes and plant carbon-water relations*, Acad. Press, Physiol. Ser., pp371-386, 1993) and to assist conventional breeding of new plant varieties in a number of species (Hall *et al.*, *Plant Breeding Reviews* 4, 81-113, 1994) including rice (Farquhar *et al.*, *In: Breaking the Yield Barrier*, ed KG Cassman, IRRI, 95, 101) and most recently wheat (Rebetzke *et al.* *Crop Science* 42:739-745, 2002).

15

No single gene has been identified as being capable of enhancing transpiration efficiency when expressed *in planta*. Transpiration efficiency may well be multigenic. As a consequence, the genes and signalling pathways that regulate the photosynthetic and/or stomatal components of the transpiration efficiency mechanism in plants have not been identified or characterized.

20

Moreover, notwithstanding that the effect of down-regulating expression of the *Rubisco* gene, or mutation in genes involved in abscissic acid (eg. *aba*, *abi*), are known to modify transpiration efficiency to some extent through stomatal closure, the consequence of such modifications is not totally specific, resulting in pleiotropic

25

effects.

30

Arabidopsis thaliana ecotype Landsberg *erecta* (L-*er*) is one of the most popular ecotypes and is used widely for both molecular and genetic studies. It harbors the *erecta* (*er*) mutation, which confers a compact inflorescence, blunt fruits, and short

petioles. There are a number of *er* mutant alleles. Phenotypic characterization of the mutant alleles suggests a role for the wild type *ER* gene in regulating plant morphogenesis, particularly the shapes of organs that originate from the shoot apical meristem. Torii *et al.*, *The Plant Cell* 8, 735, 1996, showed that the *ER* gene encodes
 5 a putative receptor protein kinase comprising a cytoplasmic protein kinase catalytic domain, a transmembrane region, and an extracellular domain consisting of leucine-rich repeats, which are thought to interact with other macromolecules.

BRIEF DESCRIPTION OF THE DRAWINGS

10 Figure 1a is a graphical representation showing the CO₂ assimilation rates ($\mu\text{mol C m}^{-2} \text{ s}^{-1}$) of several genotypes of *A. thaliana*. The genotypes of plants are indicated on the x-axis, and CO₂ assimilation rates indicated on the ordinate. Col indicates a genetic background of the ecotype Columbia. Ld indicates a genetic background of the ecotype Landsberg. Plants expressing wild type *ERECTA* alleles were either in a
 15 Col (Col4-*ER*) or Ld (Ld-*ER*) background. Plants that were homozygous for a mutant *er* allele were either in a Col background (Col-*er105* or Col-*er2*) or a Ld background (Ld-*er1*). Plants designated as F1 (Col-*ER* x Ld-*er*) were heterozygous *ER/er1*. Data indicate that, in a Col background, the *er105* mutation leads to reduced CO₂ assimilation rate, whilst the *er1* mutation enhances CO₂ assimilation rate in a Ld
 20 background.

Figure 1b is a graphical representation showing the stomatal conductance ($\text{mol H}_2\text{O m}^{-2} \text{ s}^{-1}$) of several genotypes of *A. thaliana*. The genotypes of plants are indicated on the x-axis and are the same as described in the legend to Figure 1a. Stomatal
 25 conductances are indicated on the ordinate. Data indicate that, in a Col background, the *er2* mutation significantly enhances stomatal conductance, whilst the *er1* mutation significantly enhances stomatal conductance in a Ld background.

Figure 1c is a graphical representation showing the transpiration efficiency of ($\text{mmol C mol H}_2\text{O}^{-1}$) of several genotypes of *A. thaliana*, as determined by the ratio of CO₂ assimilation rate to stomatal conductance. The genotypes of plants are indicated on the x-axis and are the same as described in the legend to Figure 1a. Transpiration
 30 efficiency is indicated on the ordinate. Data indicate that transpiration efficiency is enhanced in plants expressing a wild type *ER* allele relative to a mutant *er* allele, in

both Ld and Col backgrounds. The lowest transpiration efficiency was observed for plants that are homozygous for the *er105* allele (ie. Col-*er105*), consistent with the fact that this allele disrupts *ERECTA* expression. From the data in Figures 1a-1c, it is apparent that the lower transpiration efficiency of plants expressing the *er105* allele is largely due to a reduced CO₂ fixation rate, whereas for both the *er2* and *er1* alleles, reduced transpiration efficiency is largely due to enhanced stomatal conductance. The transpiration efficiency of the F1 heterozygote plant was intermediate between the transpiration efficiencies of its parents, suggesting codominance of these alleles. The F1, however, had a transpiration efficiency closer to that of the pollen donor parent, Ld-*er1*.

Figure 2a is a graphical representation showing the stomatal densities (Number of stomata mm⁻² leaf) for several genotypes of *A. thaliana* in three independent experiments. The genetic backgrounds of plants are indicated on the x-axis (Col, Columbia; Ld, Landsberg), and stomatal densities are indicated on the ordinate. Plant genotypes are indicated at the top of each bar, as follows: plants expressing wild type *ERECTA* alleles in a Col background were Col4*ER* or Col1*ER* (hatched bars); plants expressing wild type *ERECTA* alleles in a Ld background were *ER* (open bars); plants expressing mutant *erecta* alleles in a Col background were either *er105* or *er2* (stippled boxes); and plants expressing the mutant *er1* allele in a Ld background were *er1* (filled boxes). Columns designated a,b are data from two experiments where plants were grown in soil in the absence of fertiliser. The set of columns at the right of the figure are from a third experiment where the same plants were grown in soil comprising fertiliser. Data indicate that, in a Col background, the *er105* mutation and *er2* mutation enhances stomatal density, which in part accounts for the enhanced stomatal conductances and reduced transpiration efficiencies of plants expressing these alleles (Figures 1b and 1c). The general effects of these alleles is not dependent on the nutrient status of the soil. In contrast, the *er1* allele enhanced stomatal density of Ld plants in only one case when fertiliser was absent, suggesting that enhanced stomatal aperture may account for the enhanced stomatal conductances and reduced transpiration efficiencies of Ld-*er1* plants (Figures 1b, 1c).

Figure 2b is a graphical representation showing the epidermal cell size (surface area, μm^2) for several genotypes of *A. thaliana* in three independent experiments. The genetic backgrounds and genotypes of plants are indicated on the x-axis and at the tops of each column, respectively, as in the legend to Figure 2a. The ordinate indicates epidermal cell size. Columns designated a,b are data from two experiments where plants were grown in soil in the absence of fertiliser. The set of columns at the right of the figure are from a third experiment where the same plants were grown in soil comprising fertiliser. Data indicate that, in a Col background, the *er105* mutation and *er2* mutation significantly reduce epidermal cell size ie increase the number of epidermal cells per unit leaf area. This reveals that the ER gene has effects on leaf histogenesis which, beyond their consequences on stomatal densities, may also directly affect leaf capacity for photosynthesis and therefore transpiration efficiency, (Figures 1b and 1c). The general effects of these alleles is not dependent on the nutrient status of the soil. In contrast, the *er1* allele reduced epidermal cell size of Ld plants in only one case when fertiliser was absent, suggesting that enhanced stomatal aperture accounts for the enhanced stomatal conductances and reduced transpiration efficiencies of Ld-*er1* plants (Figures 1b, 1c).

Figure 2c is a graphical representation showing the stomatal index for several genotypes of *A. thaliana* in three independent experiments. The genetic backgrounds and genotypes of plants are indicated on the x-axis and at the tops of each column, respectively, as in the legend to Figure 2a. The ordinate indicates stomatal index, as determined from the ratio of stomatal density to epidermal cell density. Columns designated a,b are data from two experiments where plants were grown in soil in the absence of fertiliser. The set of columns at the right of the figure are from a third experiment where the same plants were grown in soil comprising fertiliser. Data indicate that the *er* mutations tested do not significantly modify stomatal index, because increases in stomatal density are correlated to increases in epidermal cell numbers in the mutant plants. Accordingly, the *ER* gene does not appear to directly modify stomatal development *per se*.

SUMMARY OF THE INVENTION

In work leading up to the present invention, the inventors sought to elucidate the specific genetic determinants of plant transpiration efficiency. In plants, the

development of molecular genetic markers, such as, for example, genetic markers that map to a region of the genome of a crop plant, such as, for example, a region of the rice genome, maize genome, barley genome, sorghum genome, or wheat genome, or a region of the tomato genome or of any Brassicaceae, assists in the production of plants having enhanced transpiration efficiency (Edwards *et al.*, *Genetics* 116, 113 - 125, 1987; Paterson *et al.*, *Nature* 335, 721-726, 1988).

The present inventors identified a locus that is linked to the genetic variation in transpiration efficiency in plants. To elucidate a locus associated with the transpiration efficiency of plants, the inventors established experimental conditions and sampling procedures to determine the contribution to total transpiration efficiency of the factors influencing this phenotype, and, more particularly, the genetic contribution to the total variation in transpiration efficiency. Factors influencing transpiration efficiency include, for example, genotype of the plant, environment (eg. temperature, light, humidity, boundary layer around the leaves, root growth conditions), development (eg. age and/or stage and/or posture of plants that modifies gas exchange and/or carbon metabolism), and seed-specific factors (Masle *et al.* 1993, *op. cit.*). The screens developed by the inventors were also used to survey mutant and wild type populations for variations in transpiration efficiency and to identify ecotypes having contrasting transpiration efficiencies including the parental lines that had been used by Lister and Dean (1993). The transpiration efficiencies of the members of Lister and Dean's (1993) Recombinant Inbred Line (RIL) mapping population were then determined, and linkage analyses were performed against genetic markers to determine the chromosome regions that are linked to genetic variation in transpiration efficiency, thereby identifying a locus conditioning transpiration efficiency.

In the exemplified embodiment of the invention, there is provided a locus associated with transpiration efficiency of *A. thaliana*, said locus defined as the *ERECTA* locus on *A. thaliana* chromosome 2. The present invention clearly extends to homologs of the *A. thaliana ERECTA* locus from other plant species, identified using the methods described herein.

Accordingly, one aspect of the invention provides a locus associated with the genetic variation in transpiration efficiency of a plant, wherein said locus comprises a nucleotide sequence linked genetically to the *A. thaliana ERECTA* locus.

- 5 As used herein, the terms "genetically linked" and "map to" shall be taken to refer to a sufficient genetic proximity between a linked nucleic acid comprising a gene, allele, marker or other nucleotide sequence and nucleic acid comprising all or part of the *A. thaliana ERECTA* locus or all or part of a homolog of the *A. thaliana ERECTA* locus from another plant species, to permit said linked nucleic acid to be useful for
10 determining the presence of a particular allele of said *A. thaliana ERECTA* locus or said homolog. Those skilled in the art will be aware that for such linked nucleic acid to be used in this manner, it must be sufficiently close to said locus not to be in linkage disequilibrium or to have a high recombination frequency between said linked nucleic acid and said locus. Preferably, the linked nucleic acid and the locus
15 are less than about 25cM apart, more preferably less than about 10cM apart, even more preferably less than about 5cM apart, still even more preferably less than about 3cM apart and still even more preferably less than about 1cM apart.

- Preferably, all or part of the locus of the invention is provided as recombinant or
20 isolated nucleic acid, such as, for example, in the form of a gene construct (eg. a recombinant plasmid or cosmid), to facilitate germplasm screening.

- The *ERECTA* locus or a gene that is linked to the *ERECTA* locus is particularly useful in a breeding program, to predict the transpiration efficiency of a plant, or
25 alternatively, as a selective breeding marker to select plants having enhanced transpiration efficiency. Once mapped, marker-assisted selection (MAS) is used to introduce the *ERECTA* locus or markers linked thereto into a wide variety of populations. MAS has the advantage of reducing the breeding population size required, and the need for continuous recurrent testing of progeny, and the time
30 required to develop a superior line.

Accordingly, a second aspect of the invention provides a method of selecting a plant having enhanced transpiration efficiency, comprising:

- (a) identifying a locus associated with genetic variation in transpiration efficiency in a plant; and
- (b) selecting a plant that comprises or expresses a gene that maps to the locus.

5 Preferably, this aspect of the invention provides a method of selecting a plant having enhanced transpiration efficiency, comprising:

- (a) identifying a locus on the Arabidopsis chromosome 2 (46-50.7 cM) associated with genetic variation in transpiration efficiency in a plant; and
- (b) selecting a plant that comprises or expresses a gene that maps to the locus.

10

An alternative embodiment provides a method of selecting a plant having enhanced transpiration efficiency, comprising selecting a plant that comprises or expresses a functionally equivalent homolog of a protein-encoding region of the *ERECTA* gene of *A. thaliana*.

15

As exemplified herein, the inventors also identified specific genes or alleles that are linked to the *ERECTA* locus and determine the transpiration efficiency of a plant. More particularly, the transpiration efficiencies of near-isogenic lines, wherein each line carries a mutation within the *ERECTA* locus were determined, thereby providing

20 the genetic contribution of genes or alleles at the *ERECTA* locus to transpiration efficiency. This analysis allowed the inventors to assess the genetic contribution of particular alleles to transpiration efficiency, thereby determining allelic variants that are linked to a particular transpiration efficiency. Thus, the elucidation of the *ERECTA* locus for transpiration efficiency in plants facilitated the fine mapping and

25 determination of allelic variants that modulate transpiration efficiency.

Accordingly, a third aspect of the invention provides a method of identifying a gene that determines the transpiration efficiency of a plant comprising:

- (a) identifying a locus associated with genetic variation in transpiration efficiency
- 30 in a plant;
- (b) identifying a gene or allele that is linked to said locus, wherein said gene or allele is a candidate gene or allele for determining the transpiration efficiency of a plant; and

- (c) determining the transpiration efficiencies of a panel of plants, wherein not all members of said panel comprise said gene or allele, and wherein variation in transpiration efficiency between the members of said panel indicates that said gene is involved in determining transpiration efficiency.

5

In another embodiment, the method comprises:

- (a) identifying a locus associated with genetic variation in transpiration efficiency in a plant;
- (b) identifying multiple alleles of a gene that is linked to said locus, wherein said gene is a candidate gene involved for determining the transpiration efficiency of a plant; and
- (c) determining the transpiration efficiencies of a panel of plants, wherein each member of said panel comprises at least one of said multiple alleles, wherein variation in transpiration efficiency between the members of said panel indicates that said gene is involved in determining transpiration efficiency.

10
15

Preferably, the identified gene or allele identified by the method described in the preceding paragraph is an *ERECTA* allele, an *erecta* allele, a homolog of *ERECTA* allele, or a homolog of *erecta* allele, wherein said homolog is from a plant species other than *A. thaliana*.

20

The identified gene or allele, including any homologs from a plant other than *A. thaliana*, such as, for example, the wild-type *ERECTA* allele or a homolog thereof, is useful for the production of novel plants. Such plants are produced, for example, using recombinant techniques, or traditional plant breeding approaches such as introgression.

25

Accordingly, a still further aspect of the present invention provides a method of enhancing the transpiration efficiency of a plant comprising ectopically expressing in a plant an isolated gene comprising a nucleotide sequence that is homologous to a protein-encoding region of a gene of *A. thaliana* that maps to the *ERECTA* locus on chromosome 2. A related embodiment of the invention provides a method of enhancing the transpiration efficiency of a plant comprising introgressing into said plant a nucleic acid comprising a nucleotide sequence that is homologous to a

30

protein-encoding region of a gene of *A. thaliana* that maps to the *ERECTA* locus on chromosome 2.

5 A further aspect of the invention provides for the use of an isolated gene comprising a nucleotide sequence that is homologous to a protein-encoding region of a gene of *A. thaliana* that maps to the *ERECTA* locus on chromosome 2 in the preparation of a gene construct for enhancing the transpiration efficiency of a plant.

10 A fifth aspect of the present invention provides a plant having enhanced transpiration efficiency, wherein said plant is produced by a method described herein.

DETAILED DESCRIPTION OF THE INVENTION

Loci for transpiration efficiency and their identification

15 One aspect of the invention provides a locus associated with the genetic variation in transpiration efficiency of a plant, wherein said locus comprises a nucleotide sequence linked genetically to the *ERECTA* locus on chromosome 2 or a homolog thereof.

20 As used herein, the term "locus" shall be taken to mean the location of one or more genes in the genome of a plant that affects a quantitative characteristic of the plant, in particular the transpiration efficiency of a plant. In the present context, a "quantitative characteristic" is a phenotype of the plant for which the phenotypic variation among different genotypes is continuous and cannot be separated into
25 discrete classes, irrespective of the number of genes that determine or control the phenotype, or the magnitude of genetic effects that single gene has in determining the phenotype, or the magnitude of genetic effects of interacting genes.

By "associated with the genetic variation in transpiration efficiency of a plant"
30 means that a locus comprises one or more genes that are expressed to determine or regulate the transpiration efficiency of a plant, irrespective of the actual rate of transpiration achieved by the plant under a specified environmental condition.

The present invention clearly contemplates the presence of multiple genes that are genetically linked or map to the specified *ERECTA* locus on chromosome 2. Without being bound by any theory or mode of action, such multiple linked genes may interact, such as, for example, by epistatic interaction, to determine the transpiration efficiency phenotype.

The present invention also contemplates the presence of different alleles of any gene that is linked to the *ERECTA* locus, wherein said allele is expressed to determine the transpiration efficiency phenotype. In one embodiment, such alleles are identified by detecting a particular transpiration efficiency phenotype that is linked to the expression of the particular allele. Alternatively, or in addition, the different alleles linked to a locus are identified by detecting a structural polymorphism in DNA (eg. a restriction fragment length polymorphism (RFLP), amplified fragment length polymorphism (AFLP), single strand chain polymorphism (SSCP), and the like), that is linked to a particular transpiration efficiency phenotype.

The present invention clearly encompasses all interacting genes and/or alleles that are genetically linked to an *ERECTA* locus and are expressed to determine a transpiration efficiency phenotype. Such linked interacting genes and/or alleles will map to a region of the genome of a plant that is homologous to a region of the *Arabidopsis thaliana* genome that is associated with the transpiration efficiency of that plant and maps to the *ERECTA* locus. Preferably, such interacting genes and/or alleles comprise a protein-encoding portion of a gene of *A. thaliana* positioned within the *ERECTA* locus of the *Arabidopsis thaliana* genome that is associated with the transpiration efficiency of that plant, or a homologous protein-encoding region from another plant species.

The terms "region of the *Arabidopsis thaliana* genome that is associated with the transpiration efficiency", and "locus of *A. thaliana* that determines the transpiration efficiency" shall be taken to mean that portion of chromosome 2 of *A. thaliana* associated with transpiration efficiency, preferably the region from about 46cM to about 50.7 cM.

Even more preferably, the locus of the invention is linked to or comprises the *ERECTA* allele or the *erecta* alleles, or a protein-encoding portion thereof or a homologous gene from another plant species.

- 5 As used herein, the term "*ERECTA*" shall be taken to refer to a wild type *A. thaliana* *ERECTA* allele or a homolog thereof from another plant species.

For the purposes of nomenclature, the nucleotide sequence of the *Arabidopsis thaliana* *ERECTA* protein-encoding region, including 5'-untranslated region (UTR)
 10 and 3'-UTR, is provided herein as SEQ ID NO: 1. The amino acid sequence of the polypeptide encoded by SEQ ID NO: 1 is set forth herein as SEQ ID NO: 2.

The term "*erecta*" shall be taken to mean any allelic variant of the wild-type *ERECTA* allele that modifies transpiration efficiency of *A. thaliana* or a homolog thereof from
 15 another plant species. Preferred *erecta* alleles contemplated herein include an *A. thaliana* *erecta* allele selected from the group consisting of: *er*, *er1*, *er2* allele, *er101* allele, *er102* allele, *er103* allele, *er104* allele, *er105* allele, *er2* allele, *er108* allele, *er109* allele, *er110* allele, *er111* allele, *er112* allele, *er113* allele, *er114* allele, *er115* allele, *er116* allele, *er117* allele, *er118* allele, *er119* allele, *er120* allele, *er121* allele,
 20 *er122* allele, *er123* allele (Lease *et al.* *New Phytologist* 151, 133-143, 2001) and a homolog of any one of said alleles.

Those skilled in the art are aware that the terms "homolog" and "ortholog" refer to functional equivalent units. In the present context, a homolog or ortholog of a gene
 25 that maps to the *ERECTA* locus shall be taken to mean any gene from a plant species other than *A. thaliana* that is functionally equivalent to a gene that maps to the exemplified *A. thaliana* *ERECTA* locus, and comprises a protein-encoding region in its native plant genome that shares a degree of structural identity or similarity with a protein-encoding region that is linked to the *A. thaliana* *ERECTA* locus.

30

Preferably, a homologous or orthologous gene from a plant other than *A. thaliana* will be associated with the transpiration efficiency of said plant and be linked to a protein-encoding region in its native plant genome that comprises a nucleotide sequence having at least about 55% overall sequence identity to a protein-encoding

region linked to the *ERECTA* locus. Even more preferably, the percentage identity will be at least about 59-61% or 70% or 80%, still more preferably at least about 90%, and even still more preferably at least about 95%.

5 In determining whether or not two nucleotide sequences fall within a particular percentage identity limitation recited herein, those skilled in the art will be aware that it is necessary to conduct a side-by-side comparison or multiple alignment of sequences. In such comparisons or alignments, differences may arise in the positioning of non-identical residues, depending upon the algorithm used to perform
10 the alignment. In the present context, reference to a percentage identity between two or more nucleotide sequences shall be taken to refer to the number of identical residues between said sequences as determined using any standard algorithm known to those skilled in the art. For example, nucleotide sequences may be aligned and their identity calculated using the BESTFIT program or other appropriate program of
15 the Computer Genetics Group, Inc., University Research Park, Madison, Wisconsin, United States of America (Devereaux *et al*, *Nucl. Acids Res.* 12, 387-395, 1984).

Alternatively, or in addition, a homologous or orthologous gene from a plant other than *A. thaliana* will be associated with the transpiration efficiency of said plant and
20 be linked to a protein-encoding region in its native plant genome that comprises a nucleotide sequence that encodes a polypeptide having at least about 55% overall sequence identity to a polypeptide encoded by a protein-encoding region linked to the *ERECTA* locus. Preferably, the percentage identity at the amino acid level will be at least about 59-61% or 70% or 80%, more preferably at least about 90%, and still
25 more preferably at least about 95%.

In determining whether or not two amino acid sequences fall within these percentage limits, those skilled in the art will be aware that it is necessary to conduct a side-by-side comparison or multiple alignment of sequences. In such comparisons
30 or alignments, differences will arise in the positioning of non-identical residues, depending upon the algorithm used to perform the alignment. In the present context, reference to a percentage identity or similarity between two or more amino acid sequences shall be taken to refer to the number of identical and similar residues respectively, between said sequences as determined using any standard algorithm

known to those skilled in the art. For example, amino acid sequence identities or similarities may be calculated using the GAP program and/or aligned using the PILEUP program of the Computer Genetics Group, Inc., University Research Park, Madison, Wisconsin, United States of America (Devereaux *et al.*, 1984, *supra*). The
5 GAP program utilizes the algorithm of Needleman and Wunsch, *J. Mol. Biol.* 48, 443-453, 1970, to maximize the number of identical/similar residues and to minimize the number and length of sequence gaps in the alignment. Alternatively or in addition, wherein more than two amino acid sequences are being compared, the ClustalW program of Thompson *et al.*, *Nucl. Acids Res.* 22, 4673-4680, 1994, is used.

10

Alternatively, or in addition, a homologous or orthologous gene from a plant other than *A. thaliana* will be associated with the transpiration efficiency of said plant and be linked to a protein-encoding region in its native plant genome that hybridizes to nucleic acid that comprises a sequence complementary to a protein-encoding region
15 linked to the *A. thaliana* *ERECTA* locus. Preferably, such homologs or orthologs will be identified by hybridization under at least low stringency conditions, and more preferably under at least moderate stringency or high stringency hybridization conditions.

20 For the purposes of defining the level of stringency, a low stringency is defined herein as being a hybridization or a wash carried out in 6xSSC buffer, 0.1% (w/v) SDS at 28°C or alternatively, as exemplified herein. Generally, the stringency is increased by reducing the concentration of salt in the hybridization or wash buffer, such as, for example, by reducing the concentration of SSC. Alternatively, or in
25 addition, the stringency is increased, by increasing the concentration of detergent (eg. SDS). Alternatively, or in addition, the stringency is increased, by increasing the temperature of the hybridization or wash. For example, a moderate stringency can be performed using 0.2xSSC to 2xSSC buffer, 0.1% (w/v) SDS, at a temperature of about 42°C to about 65°C. Similarly, a high stringency can be performed using
30 0.1xSSC to 0.2xSSC buffer, 0.1% (w/v) SDS, at a temperature of at least 55°C. Conditions for performing nucleic acid hybridization reactions, and subsequent membrane washing, are well understood by one normally skilled in the art. For the purposes of further clarification only, reference to the parameters affecting hybridization between nucleic acid molecules is found in Ausubel *et al.*, *In: Current*

Protocols in Molecular Biology, Greene/Wiley, New York USA, 1992, which is herein incorporated by reference.

Preferably, the homologous gene is derived from, or present in, the genome of a plant
 5 that is desiccation or drought intolerant, or poorly adapted for growth in dry or arid
 environments, or that suffers from reduced vigor or growth during periods of
 reduced rainfall or drought, or from the genome of a plant with increased growth
 rate or growth duration or partitioning of C to shoot and harvested parts under well-
 watered conditions.

10

More preferably, the homologous gene is derived from, or present in, the genome of a
 brassica plant, broad acre crop plant, perennial grass (eg. of the subfamily Pooidae,
 or the Tribe Poeae), or tree. Even more preferably, the homologous locus is in the
 genome of a plant selected from the group consisting of barley, wheat, rye, sorghum,
 15 rice, maize, *Phalaris aquatica*, *Dactylus glomerata*, *Lolium perenne*, *Festuca*
arundinacea, cotton, tomato, soybean, oilseed rape, poplar, and pine.

A particularly preferred homolog of a gene that maps to the exemplified *Arabidopsis*
thaliana *ERECTA* locus is derived from chromosome 6 of rice (*Oryza sativa*), and,
 20 more preferably is linked to an *ERECTA* gene derived from rice. For the purposes of
 nomenclature, the protein-encoding region of the rice *ERECTA* gene is provided
 herein as SEQ ID NO: 3. The amino acid sequence of the polypeptide encoded by
 SEQ ID NO: 3 is set forth herein as SEQ ID NO: 4.

25 Another particularly preferred homolog of a gene that maps to the exemplified
Arabidopsis thaliana *ERECTA* locus is derived from the genome of *Sorghum bicolor*,
 and, more preferably is linked to an *ERECTA* gene derived from sorghum. For the
 purposes of nomenclature, the protein-encoding region of the sorghum *ERECTA*
 gene is provided herein as SEQ ID NO: 5. The amino acid sequence of the
 30 polypeptide encoded by SEQ ID NO: 5 is set forth herein as SEQ ID NO: 6.

Another particularly preferred homolog of a gene that maps to the exemplified
Arabidopsis thaliana *ERECTA* locus is derived from the genome of *A. thaliana*, and,
 more preferably is linked to an *ERECTA* gene derived from *A. thaliana*. For the

purposes of nomenclature, the protein-encoding region of the *A. thaliana* *ERECTA* homologue is provided herein as SEQ ID NO: 7. The amino acid sequence of the polypeptide encoded by SEQ ID NO: 7 is set forth herein as SEQ ID NO: 8.

- 5 Another particularly preferred homolog of an *ERECTA* gene is provided herein as SEQ ID NO: 9. The amino acid sequence of the polypeptide encoded by SEQ ID NO: 9 is set forth herein as SEQ ID NO: 10.

A number of mapping methods for determining useful loci and estimating their
 10 effects have been described (eg. Edwards *et al.*, *Genetics* 116, 113-125, 1987; Haley and Knott, *Heredity* 69, 315-324, 1992; Jiang and Zeng, *Genetics* 140, 1111-1127, 1995; Lander and Botstein, *Genetics* 121, 185-199, 1989; Jansen and Stam, *Genetics* 136, 1447-1455, 1994; Utz and Melchinger, *In: Biometrics in Plant Breeding: Applications of Molecular Markers*. Proc. Ninth Meeting of the EUCARPIA Section
 15 Biometrics in Plant Breeding, 6 - 8 July 1994, Wageningen, The Netherlands, (J.W. van Ooijen and J. Jansen, eds), pp195-204, 1994; Zeng, *Genetics* 136, 1457-1468, 1994). In the present context, these methods are applied to identify the major component(s) of the total genetic variance that contribute(s) to the variation in transpiration efficiency of a plant, such as, for example, determined by the
 20 measurement of carbon isotope discrimination (Δ). More particularly, the segregation of known markers is used to map and/or characterize an underlying locus associated with transpiration efficiency. The locus method involves searching for associations between the segregating molecular markers and transpiration efficiency in a segregating population of plants, to identify the linkage of the marker
 25 to the locus.

To discover a marker/locus linkage, a segregating population is required. Experimental populations, such as, for example, an F₂ generation, a backcross (BC) population, recombinant inbred line (RIL), or double haploid line (DHL), can be used
 30 as a mapping population. Bulk segregant analysis, for the rapid detection of markers at specific genomic regions using segregating populations, is described by Michelmoore *et al.*, *Proc. Natl Acad. Sci. (USA)* 88, 9828-9832, 1991. In the case of F₂ mapping populations, F₂ plants are used to determine genotype, and F₂ families to determine phenotype. Recombinant inbred lines are produced by single-seed

descent. Recombinant inbred lines, such as, for example, the F9 RILs of *A. thaliana* (eg. Lister and Dean, *Plant J.*, 4, 745-750, 1993) will be known to those skilled in the art. Near isogenic lines (NILs) are used for fine mapping, and to determine the effect of a particular locus on transpiration efficiency. An advantage of recombinant inbred
5 lines and double haploid lines is that they are permanent populations, and as a consequence, provide for replication of the contribution of a particular locus to the transpiration efficiency phenotype.

As for statistical methods, Single Marker Analysis (Point Analysis) is used to detect a
10 locus in the vicinity of a single genetic marker. The mean transpiration efficiencies of a population of plants segregating for a particular marker, are compared according to the marker class. The difference between two mean transpiration efficiencies provides an estimate of the phenotypic effect of substituting one allele for another allele at the locus. To determine whether or not the inferred phenotypic effect is
15 significantly different from zero, a simple statistical test, such as t-test or F-test, is used. A significant value indicates that a locus is located in the vicinity of the marker. Single point analysis does not require a complete molecular linkage map. The further locus is from the marker, the less likely it is to be detected statistically, as a consequence of recombination between the marker and the gene.

20

In the Anova, t-test or GLM approach, the association between marker genotype and transpiration efficiency phenotype comprises:

- (i) classifying progeny of a segregating population of plants by marker genotype, such as for example, using RFLP, AFLP, SSCP, or microsatellite analyses,
25 thereby establishing classes of plants;
- (ii) comparing the mean transpiration efficiencies of classes of plants in the segregating population, using a t-test, GLM or ANOVA; and
- (iii) determining the significance of the differences in the mean at (ii), wherein a significant difference indicates that the marker is linked to the locus for
30 transpiration efficiency.

As will be known to those skilled in the art, the difference between the means of the classes provides an estimate of the effect of the locus in determining the transpiration efficiency of a class.

In the regression approach, the association between marker genotype and phenotype is determined by a process comprising:

- (i) assigning numeric codes to marker genotypes; and
 - 5 (ii) determining the regression value r for transpiration efficiency against the codes, wherein a significant value for r indicates that the marker is linked to the locus for transpiration efficiency, and wherein the regression slope estimate of the effect of a particular locus on transpiration efficiency.
- 10 For QTL interval mapping, the Mapmaker algorithm developed by Lincoln *et al.*, Constructing genetic linkage maps with MAPMAKER/EXP version 3.0: A tutorial and reference manual. Whitehead Institute for Biomedical Research, Cambridge, MA, USA, 1993, can be used. The principle behind interval mapping is to test a model for the presence of a QTL at many positions between two mapped marker loci. This
- 15 model is a fit of a presumptive QTL to transpiration efficiency, wherein the suitability of the fit is tested by determining the maximum likelihood that a QTL, for transpiration efficiency lies between two segregating markers. For example, in the case of a QTL located between two segregating markers, the 2-loci marker genotypes of segregating progeny will each contain mixtures of QTL genotypes. Accordingly, it
- 20 is possible to search for loci parameters that best approximate the distribution in transpiration efficiency for each marker class. Models are evaluated by computing the likelihood of the observed distributions with and without fitting a QTL effect. The map position of a QTL is determined as the maximum likelihood from the distribution of likelihood values (LOD scores: ratio of likelihood that the effect
- 25 occurs by linkage: likelihood that the effect occurs by chance), calculated for each locus.

Interval mapping by regression (Haley and Knott., *Heredity* 69, 315-324, 1992) is a simplification of the maximum likelihood method *supra* wherein basic QTL analysis

30 or regression on coded marker genotypes is performed, except that phenotypes are regressed on the probability of a QTL genotype as determined from the linkage between transpiration efficiency and the nearest flanking markers. In most cases, regression mapping gives estimates of QTL position and effect that are almost

identical to those given by the maximum likelihood method. The approximation deviates only at places where there are large gaps, or many missing genotypes.

In the composite interval mapping (CIM) method (Jansen and Stam, *Genetics* 136, 1447-1455, 1994; Utz and Melchinger, 1994, *supra*; Zeng, *Genetics* 136, 1457-1468, 1994), the analysis is performed in the usual way, except that the variance from other QTLs are accounted for by including partial regression more power and precision than simple interval mapping, because the effects of other QTLs are not present as residual variance. CIM can remove the bias that can be caused by the QTLs that are linked to the position being tested.

Publicly available software are used to map a locus for transpiration efficiency. Such software include, for example, the following:

- (i) MapMaker/QTL (<ftp://genome.wi.mit.edu/pub/mapmaker3/>), for analyzing F2 or backcross data using standard interval mapping;
- (ii) MQTL, for composite interval mapping in multiple environments or for performing simple interval mapping using homozygous progeny (eg. double haploids, or recombinant inbred lines);
- (iii) PLABQTL (Utz and Melchinger, PLABlocus Version 1.0. A computer program to map QTL, Institut für Pflanzenzüchtung, Saatgutforschung und Populationsgenetik, Universität Hohenheim, 70593 Stuttgart, Germany, 1995; <http://www.uni-hohenheim.de/~ipspwww/soft.html>) for composite interval mapping and simple interval mapping of a locus in mapping populations derived from a bi-parental cross by selfing, or in double haploids;
- (iv) QTL Cartographer (<http://statgen.mcsu.edu/qtlcart/cartographer.html>) for single-marker regression, interval mapping, or composite interval mapping, using F2 or backcross populations;
- (v) MapQTL (<http://www.cpro.dlo.nl/cbw/>); Qgene for performing either single-marker regression or interval regression to map loci; and
- (vi) SAS for detecting a locus by identifying associations between marker genotype and transpiration efficiency by a single marker analysis approach such as ANOVA, t-test, GLM or REG.

In a particularly preferred embodiment, QTL cartographer or MQTL is used to identify a locus associated with the transpiration efficiency of plants.

Those skilled in the art will also be aware that it is possible to detect multiple
5 interacting alleles or genes for a particular trait, such as, for example, using
composite interval mapping approaches. To achieve this end, the composite interval
mapping may be repeated to look for additional loci. Alternatively, or in addition,
two or more distinct regions of the genome can be nominated as candidate loci, and
a gamete relationship matrix constructed for each candidate locus, and a 2-locus
10 regression performed for each pair of loci, determining a best fit for the interacting
effects between the two loci or alleles at those loci, including any dominance or
additive effects. The algorithm described by Carlborg *et al.*, *Genetics* (2000) can be
used for simultaneous mapping. In the present context, such an analysis is
performed with reference to the segregation of transpiration efficiency phenotypes in
15 the segregating population.

Use of the ERECTA locus to enhance transpiration efficiency of plants

As will be known to those skilled in the art, a single locus, if present in the genome
of a plant, can have a significant influence on the phenotype of the plant. For
20 example, Grandillo *et al.*, *Theor. Appl. Genet.* 99, 978-987, 1999, showed that for
tomato a selection made from a total 28 loci determining fruit size and weight
explained 20% of the total phenotypic variance in this trait.

Accordingly, a second aspect of the invention provides a method of selecting a plant
25 having enhanced transpiration efficiency, comprising:

- (a) identifying a locus associated with genetic variation in transpiration efficiency in
a plant; and
- (b) selecting a plant that comprises or expresses a gene that maps to the locus.

30 By "enhanced transpiration efficiency" is meant that the plant loses less water per
unit of dry matter produced, or alternatively, produces an enhanced amount of dry
matter per unit of water transpired, relative to a counterpart plant. By "counterpart
plant" is meant a plant having a similar or near-identical genetic background, such
as, for example, a near-isogenic plant, a sibling, or parent.

In accordance with this aspect of the invention, a locus is identified by conventional locus mapping means, and/or by homology searching for genes that map to the *ERECTA* locus on chromosome 2 of the *A. thaliana* genome, such as, for example, by
5 searching for homologs of the *A. thaliana* *ERECTA* allele or *erecta* allele as described herein above.

Preferably, to select a plant that comprises or expresses the appropriate gene, marker-assisted selection (MAS) is used. As will be known to those skilled in the
10 art, once a particular locus has been identified, genetic or physical markers that are linked to the locus can be readily identified and used to confirm the presence of the locus in breeding populations. For a locus that is flanked by two tightly-linked markers that recombine only at a low frequency, the presence of the flanking markers is indicative of the presence of the locus.

15 For marker-assisted selection, it is preferred that the marker is a genetic marker (eg. a gene or allele), or a physical marker (eg. leaf hairiness or pod shape), or a molecular marker such as, for example, a restriction fragment length polymorphism (RFLP), a restriction (RAPD), amplified fragment length polymorphism (AFLP), or a short
20 sequence repeat (SSR) such as a microsatellite, or SNP. It is also within the scope of the invention to utilize any hybridization probe or amplification primer comprising at least about 10 nucleotides in length derived from a chromosome region that is linked in the genome of a plant to the *ERECTA* locus, as a marker to select plants. Those skilled in the art will readily be able to determine such probes or primers
25 based upon the disclosure herein, particularly for those plant genomes which may have sufficient chromosome sequence in the region of interest in the genome (eg. *A. thaliana*, rice, cotton, barley, wheat, sorghum, maize, tomato, etc).

For flanking markers that are not tightly linked, such that there is a large
30 recombination distance there between, the presence of the appropriate gene is assessed by identifying those plants having both flanking markers and then selecting from those plants a plant having an enhanced transpiration efficiency. Naturally, the greater the distance between two markers, the larger the population of plants required to identify a plant having both markers, the intervening locus and a gene

within said locus. Those skilled in the art will readily be able to determine the population size required to identify a plant having a particular transpiration efficiency, based upon the recombination units (cM) between two markers.

- 5 Transpiration efficiency is determined by any means known to the skilled artisan. Preferably, transpiration efficiency is determined by measuring dry matter accumulation in the plant by gravimetric means, or by measuring water loss, or the ratio of CO₂ assimilation rate to stomatal conductance.
- 10 In a particularly preferred embodiment, the transpiration efficiency is determined directly, by measuring the ratio of carbon fixed carbon assimilation rate) to water loss (transpiration rate).

In an alternative embodiment, transpiration efficiency is determined indirectly from
 15 the carbon isotope discrimination value (Δ). Farquhar *et al.*, *Aust. J. Plant Physiol.* 9,121-137, 1982, showed that carbon isotope discrimination (Δ ; a measure of the extent to which the ¹³C/¹²C ratio of organic matter is less than that of CO₂ in the source air), is an effective indirect measure of transpiration efficiency. The determination of transpiration efficiency in this manner is based upon the constancy
 20 of the atmospheric ¹³C: ¹²C ratio (about 98.19: 1.11) and the finding that, in C₃ plants at least, ribulose biphosphate carboxylase (Rubisco) enzymes discriminate against the use of ¹³C. Thus, ¹³CO₂ is less efficiently assimilated than ¹²CO₂, and diffuses less through stomata in and out of the leaf. However, when the stomata become nearly closed, the diffusion of ¹³CO₂ is more difficult to achieve and, at higher
 25 intracellular concentrations of ¹³CO₂, this isotope is incorporated into 3-phosphoglycerate, and subsequently into dry matter. As a consequence, carbon isotope discrimination (Δ) is greatest when the overall CO₂ assimilation rate during photosynthesis (A) is small, and stomatal conductance (g_w) to water vapor is large. This relationship is represented by the following algorithm:

$$30 \quad \Delta (\text{‰}) = 27 - 36A / (G_w \times C_a)$$

wherein C_a is the ambient CO₂ concentration (ie. [¹²CO₂ + ¹³CO₂]).

For a C₃ plant that exhibits a value in the range of about 4.5 ‰ to about 6.7 ‰ for the term 36A/(G_w × C_a), a 1 ‰ change in carbon isotope discrimination (Δ)

corresponds to a change in transpiration efficiency in the range of about 22% to about 15%, respectively.

The negative relationship between carbon isotope discrimination (Δ) and transpiration efficiency has been established for many plant species, including wheat (Farquhar and Richards, *Aust. J. Plant Physiol.* 11, 539-552, 1984; Farquhar *et al.*, *Ann. Rev. Plant Physiol.* 40, 388-397, 1989), *Stylosanthes* (Thumma *et al.*, *Proc. 9th Aust. Agronomy Conf.*, Wagga Wagga New South Wales, Australia, 1998), cotton, barley, and rice. Accordingly, a lower carbon isotope discrimination (Δ) value for a test plant relative to a counterpart plant is indicative of enhanced transpiration efficiency.

Alternatively, or in addition, transpiration efficiency is determined by another indicator, such as, for example, leaf temperature, ash content, mineral content, or specific leaf weight (dry matter per unit leaf area). For example, specific leaf weight is positively correlated with transpiration efficiency in peanuts and other species (Virgona *et al.*, *Aust. J. Plant Physiol.*, 17, 207-214, 1990; Wright *et al.*, *Crop Sci* 34, 92-97, 1994). Accordingly, a higher specific leaf weight or higher carbon gain rate for a test plant relative to a counterpart plant is indicative of enhanced transpiration efficiency of the test plant.

The presence of the locus can be established by hybridizing a probe or primer that is linked to an *ERECTA* locus, such as, for example, a probe or primer that hybridizes to the identified chromosome 2 region of *A. thaliana* or the identified chromosome 6 region of rice.

Preferably, the presence of the locus is established by hybridizing a probe or primer derived from any one or more of SEQ ID Nos: 1, 3, or 5, or from a homologous gene in another plant, or a complementary sequence to such a sequence, to genomic DNA from the plant, and detecting the hybridization using a detection means.

In one embodiment, detection of the hybridization is performed preferably by labelling a probe with a reporter molecule capable of producing an identifiable signal, prior to hybridization, and then detecting the signal after hybridization.

Preferred reporter molecules include radioactively-labelled nucleotide triphosphates and biotinylated molecules. Preferably, variants of the genes exemplified herein, including genomic equivalents, are isolated by hybridisation under moderate stringency or more preferably, under high stringency conditions, to the probe.

5

Alternatively, or in addition, hybridization may be detected using any format of the polymerase chain reaction (PCR), including AFLP. For PCR, two non-complementary nucleic acid primer molecules comprising at least about 20 nucleotides in length, and more preferably at least 30 nucleotides in length are hybridized to different
10 strands of a nucleic acid template molecule, and specific nucleic acid molecule copies of the template are amplified enzymatically. Several formats of PCR are described in McPherson *et al.*, In: *PCR A Practical Approach.*, IRL Press, Oxford University Press, Oxford, United Kingdom, 1991, which is incorporated herein by reference.

15

For enhancing the transpiration efficiency of a plant wherein the locus is polymorphic, such as, for example, an allele, the method *supra* is modified to include the detection of the specific allele(s) linked to the desired enhancement. According to this embodiment, there is provided a method of selecting a plant
20 having enhanced transpiration efficiency, comprising:

- (a) identifying a locus associated with genetic variation in transpiration efficiency in a plant;
- (b) identifying a polymorphic marker within said locus that is linked to enhanced transpiration efficiency; and
- 25 (c) selecting a plant that comprises or expresses the marker.

Standard means known to the skilled artisan are used to identify a marker within the locus that is linked to enhanced transpiration efficiency. A population of plants that is segregating for the polymorphic marker is generally used, wherein the
30 transpiration efficiency phenotype of plants is then correlated or associated with the presence of a particular allelic form of the marker. As exemplified herein, near-isogenic or recombinant inbred lines of plants were screened to segregate alleles at the *ERECTA* locus and to correlate enhanced transpiration efficiency with the presence of the *ERECTA* allele as opposed to an *erecta* allele.

Suitable markers include any one or more of the markers described herein to be suitable for MAS.

- 5 Preferably, the selection of plants in accordance with these embodiment includes the additional step of introducing the locus or polymorphic marker to a plant, such as, for example, by standard breeding approaches or by recombinant means. This may be carried out at the same time, or before, selecting the locus or polymorphic marker.
- 10 Recombinant means generally include introducing a gene construct comprising the locus or marker into a plant cell, selecting transformed tissue and regenerating a whole plant from the transformed tissue explant. Means for introducing recombinant DNA into plant tissue or cells include, but are not limited to, transformation using CaCl_2 and variations thereof, in particular the method described by Hanahan (1983),
15 direct DNA uptake into protoplasts (Krens *et al*, *Nature* 296, 72-74, 1982; Paszkowski *et al.*, *EMBO J.* 3, 2717-2722, 1984), PEG-mediated uptake to protoplasts (Armstrong *et al.*, *Plant Cell Rep.* 9, 335-339, 1990) microparticle bombardment, electroporation (Fromm *et al.*, *Proc. Natl. Acad. Sci. (USA)*, 82, 5824-5828, 1985), microinjection of DNA (Crossway *et al.*, *Mol. Gen. Genet.* 202, 179-185, 1986), microparticle
20 bombardment of tissue explants or cells (Christou *et al*, *Plant Physiol.* 87, 671-674, 1988; Sanford, *Part. Sci. Technol.* 5, 27-37, 1988), vacuum-infiltration of tissue with nucleic acid, or in the case of plants, T-DNA-mediated transfer from *Agrobacterium* to the plant tissue as described essentially by An *et al.*, *EMBO J.* 4, 277-284, 1985; Herrera-Estrella *et al.*, Herrera-Estella *et al.*, *Nature* 303, 209-213, 1983; Herrera-
25 Estella *et al.*, *EMBO J.* 2, 987-995, 1983; or Herrera-Estella *et al.*, *In: Plant Genetic Engineering*, Cambridge University Press, N.Y., pp 63-93, 1985..

- For microparticle bombardment of cells, a microparticle is propelled into a cell to produce a transformed cell. Any suitable ballistic cell transformation methodology
30 and apparatus can be used in performing the present invention. Exemplary apparatus and procedures are disclosed by Stomp *et al.* (U.S. Patent No. 5,122,466) and Sanford and Wolf (U.S. Patent No. 4,945,050). When using ballistic transformation procedures, the gene construct may incorporate a plasmid capable of replicating in the cell to be transformed.

Examples of microparticles suitable for use in such systems include 1 to 5 micron gold spheres. The DNA construct may be deposited on the microparticle by any suitable technique, such as by precipitation.

5

A whole plant may be regenerated from the transformed or transfected cell, in accordance with procedures well known in the art. Plant tissue capable of subsequent clonal propagation, whether by organogenesis or embryogenesis, may be transformed with a gene construct of the present invention and a whole plant
10 regenerated therefrom. The particular tissue chosen will vary depending on the clonal propagation systems available for, and best suited to, the particular species being transformed. Exemplary tissue targets include leaf disks, pollen, embryos, cotyledons, hypocotyls, megagametophytes, callus tissue, existing meristematic tissue (eg., apical meristem, axillary buds, and root meristems), and induced
15 meristem tissue (eg., cotyledon meristem and hypocotyl meristem).

The term "organogenesis", as used herein, means a process by which shoots and roots are developed sequentially from meristematic centres.

20 The term "embryogenesis", as used herein, means a process by which shoots and roots develop together in a concerted fashion (not sequentially), whether from somatic cells or gametes.

The generated transformed plants may be propagated by a variety of means, such as
25 by clonal propagation or classical breeding techniques. For example, a first generation (or T1) transformed plant may be selfed to give homozygous second generation (or T2) transformant, and the T2 plants further propagated through classical breeding techniques.

30 The generated transformed organisms contemplated herein may take a variety of forms. For example, they may be chimeras of transformed cells and non-transformed

cells; clonal transformants (eg., all cells transformed to contain the expression cassette); grafts of transformed and untransformed tissues (eg., in plants, a transformed root stock grafted to an untransformed scion).

5 Alternatively, the transformed plants are produced by an *in planta* transformation method using *Agrobacterium tumefaciens*, such as, for example, the method described by Bechtold *et al.*, *CR Acad. Sci. (Paris, Sciences de la vie/ Life Sciences)* 316, 1194-1199, 1993 or Clough *et al.*, *Plant J* 16: 735-74, 1998, wherein *A. tumefaciens* is applied to the outside of the developing flower bud and the binary
10 vector DNA is then introduced to the developing microspore and/or macrospore and/or the developing seed, so as to produce a transformed seed. Those skilled in the art will be aware that the selection of tissue for use in such a procedure may vary, however it is preferable generally to use plant material at the zygote formation stage for *in planta* transformation procedures.

15

Identification of genes for determining the transpiration efficiency of a plant

As exemplified herein, the inventors also identified specific genes or alleles that are linked to the *ERECTA* locus and determine the transpiration efficiency of a plant. More particularly, the transpiration efficiencies of near-isogenic lines, wherein each
20 line carries a mutation within a target locus in the region of a locus associated with transpiration efficiency, were determined, thereby providing the genetic contribution of that locus to transpiration efficiency. This analysis allowed the inventors to assess the genetic contribution of particular alleles to transpiration efficiency, thereby determining allelic variants that are linked to a particular transpiration efficiency.
25 Thus, the elucidation of the *ERECTA* locus for transpiration efficiency in plants facilitated the fine mapping and determination of allelic variants that determine transpiration efficiency.

Accordingly, a third aspect of the invention provides a method of identifying a gene
30 that determines the transpiration efficiency of a plant.

In one embodiment, the method comprises:

- (a) identifying a locus associated with genetic variation in transpiration efficiency in a plant;
- (b) identifying a gene or allele that is linked to said locus, wherein said gene or allele is a candidate gene or allele for determining the transpiration efficiency of a plant; and
- (c) determining the transpiration efficiencies of a panel of near isogenic plants, wherein not all members of said panel comprise said gene or allele, and wherein variation in transpiration efficiency between the members of said panel indicates that said gene is involved in determining transpiration efficiency.

In another embodiment, the method comprises:

- (a) identifying a locus associated with genetic variation in transpiration efficiency in a plant;
- (b) identifying multiple alleles of a gene that is linked to said locus, wherein said gene is a candidate gene involved for determining the transpiration efficiency of a plant; and
- (c) determining the transpiration efficiencies of a panel of near isogenic plants, wherein each member of said panel comprises at least one of said multiple alleles, wherein variation in transpiration efficiency between the members of said panel indicates that said gene is involved in determining transpiration efficiency.

In the present context, the term "near isogenic plants" shall be taken to mean a population of plants having identity over a substantial proportion of their genomes, notwithstanding the presence of sufficiently few differences to permit the contribution of a distinct allele or gene to the transpiration efficiency of a plant to be determined by a comparison of the transpiration efficiency phenotypes of the population. As will be known to the skilled artisan, recombinant inbred lines, lines produced by introgression of a gene followed by several generations of backcrossing, or siblings, are suitable near-isogenic lines for the present purpose.

Preferably, the identified gene or allele identified by the method described in the preceding paragraph is selected from the group consisting of *ERECTA* gene, *Erecta*

alleles, homologs of *ERECTA*, , wherein said homologs are from plants species other than *A. thaliana*.

In a particularly preferred embodiment, the identified gene or allele will comprise a nucleotide sequence selected from the group consisting of:

- (a) a sequence having at least about 55% identity to a sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, and SEQ ID NO: 9;
- (b) a sequence encoding an amino acid sequence having at least about 55% identity to an amino acid sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8 and SEQ ID NO: 10; and
- (c) a sequence complementary to (a) or (b).

Preferably, the percentage identity is at least about 59-61% or 70% or 80%, more preferably at least about 90%, and even more preferably at least about 95% or 99%. In a particularly preferred embodiment, the identified gene or allele comprises a nucleotide sequence selected from the group consisting of:

- (a) a sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, and SEQ ID NO: 9;
- (b) a sequence encoding an amino acid sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8 and SEQ ID NO: 10; and
- (c) a sequence complementary to (a) or (b).

25

Enhancement of transpiration efficiency using isolated genes

The identified gene or alleles, including any homologs from a plant other than *A. thaliana*, such as, for example, the wild-type *ERECTA* allele, or a homolog thereof, is useful for the production of novel plants. Such plants are produced, for example, using recombinant techniques, or traditional plant breeding approaches such as by introgression.

Accordingly, a fourth aspect of the present invention provides a method of enhancing the transpiration efficiency of a plant comprising ectopically expressing

in a plant an isolated gene comprising a nucleotide sequence that is homologous to a protein-encoding region of a gene that is linked to the *A. thaliana* *ERECTA* locus on chromosome 2.

5 In a particularly preferred embodiment, the isolated gene comprises a nucleotide sequence selected from the group consisting of:

- (a) a sequence having at least about 55% identity to a sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, and SEQ ID NO: 9;
- 10 (b) a sequence encoding an amino acid sequence having at least about 55% identity to an amino acid sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8 and SEQ ID NO: 10; and
- (c) a sequence complementary to (a) or (b).

15

Preferably, the percentage identity is at least about 59-61% or 70% or 80%, more preferably at least about 90%, and even more preferably at least about 95% or 99%.

In a particularly preferred embodiment, the isolated gene or allele comprises a
20 nucleotide sequence selected from the group consisting of:

- (a) a sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, and SEQ ID NO: 9;
- (b) a sequence encoding an amino acid sequence selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8 and
25 SEQ ID NO: 10; and
- (c) a sequence complementary to (a) or (b).

To ectopically express the isolated gene in a plant, the protein-encoding portion of the gene is generally placed in operable connection with a promoter sequence that is
30 operable in the plant, which may be the endogenous promoter or alternatively, a heterologous promoter, and a transcription termination sequence, which also may be the endogenous or an heterologous sequence relative to the gene of interest. The promoter and protein-encoding portion and transcription termination sequence are generally provided in the form of a gene construct, to facilitate introduction and

maintenance of the gene in a plant where it is to be ectopically expressed. Numerous vectors suitable for introducing genes into plants have been described and are readily available. These may be adapted for expressing an isolated gene in a plant to enhance transpiration efficiency therein.

5

Reference herein to a "promoter" is to be taken in its broadest context and includes the transcriptional regulatory sequences of a classical eukaryotic genomic gene, including the TATA box which is required for accurate transcription initiation, with or without a CCAAT box sequence and additional regulatory elements (ie. upstream
10 activating sequences, enhancers and silencers) which alter gene expression in response to developmental and/or external stimuli, or in a tissue-specific manner. In the present context, the term "promoter" is also used to describe a synthetic or fusion molecule, or derivative which confers, activates or enhances expression of said sense molecule in a cell. Preferred promoters may contain additional copies of one or
15 more specific regulatory elements, to further enhance expression and/or to alter the spatial expression and/or temporal expression of a nucleic acid molecule to which it is operably connected. For example, copper-responsive regulatory elements may be placed adjacent to a heterologous promoter sequence driving expression of a nucleic acid molecule to confer copper inducible expression thereon.

20

Placing a nucleic acid molecule under the regulatory control of a promoter sequence means positioning said molecule such that expression is controlled by the promoter sequence. A promoter is usually, but not necessarily, positioned upstream or 5' of the protein-encoding portion of the gene that it regulates. Furthermore, the
25 regulatory elements comprising a promoter are usually positioned within 2 kb of the start site of transcription of the structural protein-encoding nucleotide sequences, or a chimeric gene comprising same. In the construction of heterologous promoter/structural gene combinations it is generally preferred to position the promoter at a distance from the gene transcription start site that is approximately the
30 same as the distance between that promoter and the gene it controls in its natural setting, ie., the gene from which the promoter is derived. As is known in the art, some variation in this distance can be accommodated without loss of promoter function. Similarly, the preferred positioning of a regulatory sequence element with respect to a heterologous gene to be placed under its control is defined by the

positioning of the element in its natural setting, ie., the genes from which it is derived. Again, as is known in the art, some variation in this distance can also occur.

- 5 Promoters suitable for use in gene constructs of the present invention include those promoters derived from the genes of viruses, yeasts, moulds, bacteria, insects, birds, mammals and plants which are capable of functioning in plant cells, including monocotyledonous or dicotyledonous plants, or tissues or organs derived from such cells. The promoter may regulate gene expression constitutively, or differentially
10 with respect to the tissue in which expression occurs or, with respect to the developmental stage at which expression occurs, or in response to external stimuli such as physiological stresses, pathogens, or metal ions, amongst others.

Examples of promoters useful in performing this embodiment include the CaMV 35S
15 promoter, NOS promoter, octopine synthase (OCS) promoter, *Arabidopsis thaliana* SSU gene promoter, napin seed-specific promoter, SCSV promoter, SCBV promoter and the like. In addition to the specific promoters identified herein, cellular promoters for so-called housekeeping genes, including the actin promoters, or promoters of histone-encoding genes, are useful.

20

The term "terminator" refers to a DNA sequence at the end of a transcriptional unit which signals termination of transcription. Terminators are 3'-non-translated DNA sequences containing a polyadenylation signal, that facilitate the addition of a polyadenylate sequence to the 3'-end of a primary transcript. Terminators active in
25 cells derived from viruses, yeasts, moulds, bacteria, insects, birds, mammals and plants are known and described in the literature. They are isolatable from bacteria, fungi, viruses, animals and/or plants.

Examples of terminators particularly suitable for use in the gene constructs of the
30 present invention include the nopaline synthase (NOS) gene terminator of *Agrobacterium tumefaciens*, the terminator of the Cauliflower mosaic virus (CaMV) 35S gene, the *zein* gene terminator from *Zea mays*, the Rubisco small subunit (SSU) gene terminator sequences and subclover stunt virus (SCSV) gene sequence terminators, amongst others.

Those skilled in the art will be aware of additional promoter sequences and terminator sequences that may be suitable for use in performing the invention. Such sequences may readily be used without any undue experimentation.

5

Preferably, the gene construct further comprises an origin of replication sequence for its replication in a specific cell type, for example a bacterial cell, when said gene construct is required to be maintained as an episomal genetic element (eg. plasmid or cosmid molecule) in said cell. Preferred origins of replication include, but are not
10 limited to, the *f1*-ori and *colE1* origins of replication.

Preferably, the gene construct further comprises a selectable marker gene or genes that are functional in a cell into which said gene construct is introduced.

15 As used herein, the term "selectable marker gene" includes any gene which confers a phenotype on a cell in which it is expressed to facilitate the identification and/or selection of cells which are transfected or transformed with a gene construct of the invention or a derivative thereof.

20 Suitable selectable marker genes contemplated herein include the ampicillin resistance (*Amp^r*), tetracyclin-resistance gene (*Tc^r*), bacterial kanamycin resistance gene (*Kan^r*), phosphinothricin resistance gene, neomycin phosphotransferase gene (*np^tII*), hygromycin resistance gene, β -glucuronidase (GUS) gene, chloramphenicol acetyltransferase (CAT) gene, and luciferase gene, amongst others.

25

In a related embodiment, the invention extends to the use of an isolated gene comprising a nucleotide sequence that is homologous to a protein-encoding region of a gene of *A. thaliana* that is positioned between about 46cM to about 50.74cM on chromosome 2 in the preparation of a gene construct for enhancing the transpiration
30 efficiency of a plant.

In an alternative embodiment of the invention, the transpiration efficiency of a plant is enhanced by classical breeding approaches, comprising introgressing the isolated gene into a plant. For introgression of a gene, the gene is transferred from its native

genetic background into another genetic background using standard breeding, for example, a gene that enhances transpiration efficiency in a progenitor such as a diploid cotton or diploid wheat may be transferred into a commercial tetraploid cotton or hexaploid wheat, respectively, by standard crossing, followed by several
 5 generations of back-crossing to remove the genetic background of the progenitor. Naturally, continued selection of the gene of interest is required, such as, for example, facilitated by the use of markers.

A fifth aspect of the present invention provides a plant having enhanced
 10 transpiration efficiency, wherein said plant is produced by a method described herein.

The present invention is further described with reference to the following non-limiting examples.

15

EXAMPLE 1

$^{12}\text{C}/^{13}\text{C}$ discrimination as a marker for screening genetic variation
 in transpiration efficiency.

20 Experimental conditions and sampling procedures were established to allow the control of many factors, other than genetic, that influence transpiration efficiency at the level of individual leaves and plants. These factors fall into several categories: (a) characteristics of the seedling's micro-environment: temperature, light, humidity, boundary layer around the leaves, root growth conditions; (b) developmental and
 25 morphological effects that modify gas exchange and C metabolism and therefore carbon isotopic signature (eg age, stage, posture); and (c) seed effects.

We developed high resolution mass-spectrometer techniques for measuring C isotope ratios in whole tissues or carbon compounds such as soluble sugars -ie a measure of
 30 integrated transpiration efficiency over the plant's life or over a day, respectively, and also for measuring instantaneous transpiration efficiency during gas exchange.

This means:

- 0.1 per mil analytical precision in the measurement of the isotopic composition of leaf carbon. Discrimination, (δ), is approximately the isotope ratio of carbon in source CO_2 minus that of plant organic carbon. In a particular experiment, the source CO_2 is common to all genotypes.
- 5 • 0.1 per mil biological precision, that is variation between replicated seedlings, grown in soil, either in growth chambers or in glasshouses with CO_2 , humidity and temperature control (corresponding to approximately 1.5% variation in transpiration efficiency).
- The ability to grow and screen large batches of seedlings in glasshouses or growth chambers (up to 1500), under standardised leaf and root growth conditions, to a rosette size of several cm within 2-3 weeks allowing individual measurements, on the same plant, of isotope ratios and also of the underlying properties (eg in situ measurement of leaf temperature by infra-red thermometry as a measure of stomatal conductance; chlorophyll fluorescence; leaf expansion).
- 15

EXAMPLE 2

Natural genetic variation in transpiration efficiency in *Arabidopsis thaliana*

A. thaliana ecotypes were screened for leaf Δ under glasshouse conditions. There was a large spread of values (corresponding to approximately 30 % genetic variation in transpiration efficiency). However, large environmental effects were noted. A few contrasted ecotypes were selected at the two extremes of the range of Δ values and compared under various conditions of irradiance (150 to 500 $\mu\text{E m}^{-2}\text{s}^{-1}$), light spectrum (Red/Far-Red ratios) and air humidity (60 to 90%) while roots were always well watered. The magnitude of genetic differences in transpiration efficiency was very much influenced by environmental conditions. This was in part due to variations among ecotypes in the dependence of photosynthesis on light and vapour pressure deficit. Genetic differences were maximal under a combination of high light and low humidity, in growth chambers.

30

The ecotypes *Columbia (Col)* and *Landsberg erecta (Ld-er)* have extreme carbon isotope discrimination values, with *Col* always having smaller values than *Ld-er*, and thus a greater transpiration efficiency.

EXAMPLE 3

Identification of a locus associated with transpiration efficiency in *A. thaliana*
Quantitative Trait Loci (QTL) analysis of the Lister and Dean's (1993) Recombinant
Inbred Lines (later referred to as RILs) was performed to identify and map a locus
5 associated with carbon isotope discrimination (Δ). The RILs were from a cross
between Col-4 and Ler-0. Our analysis confirmed the importance of genes around
the *ER* locus, and a role for genes other than *ERECTA* in conferring transpiration
efficiency on *A. thaliana*.

10 More particularly, 300 RI mapping lines between Col and Ler ecotypes, available at
the Arabidopsis Stock Centre, were generated from a cross between the *Arabidopsis*
ecotypes Columbia (Col4) and Landsberg erecta (Ler-0 carrying *er1*) (Lister and Dean,
1993), using Columbia as the male parent. A subset of 100 of these lines, chosen as
the most densely and reliably mapped were used in the present analysis.

15

The seeds were multiplied in a glasshouse in an attempt to minimize confounding
seed effects in our comparisons. Large numbers of seeds were obtained for most lines
except for a few, including Col4 parent, which had to be re-ordered following low
seed viability of the original sample sent by the Stock Centre. The seeds harvested in
20 these propagation runs were used throughout all our experiments to date.

Loci were analysed using two programs, QTL cartographer and MQTL. These
programs compute statistics of a trait at each marker position, using a range of
methods [linear regression (LR), stepwise regression (SR), and likelihood approaches
25 (Single interval mapping (SIM) which treats values at individual markers as
independent values, and composite interval mapping (CIM) which allows for
interactions between markers and associated locus)]. By nature each of these
methods has some biases and embedded assumptions, hence the importance of
analysing data with more than one program. Only results that were consistent
30 between the two programs, and robust to additions or deletions to the set of
background markers used for composite interval mapping are reported below.

Initial QTL analysis was done in parallel to seed multiplication on a subset of 40
lines for which enough seeds were sent. Once all seeds had been multiplied this was

repeated on the full set of 100 lines. These two analyses indicated the existence of a locus for carbon isotope discrimination (Δ), that maps to the region including the *ERECTA* locus on chromosome 2, at approximately 46-51 cM (Table 1, run 1&2).

- 5 Given the complexity and integrative nature of Δ as a physiological trait, such a small number of loci associated with the trait was not expected. Subsequent experiments were therefore designed to test these results and assess their stability across the range of environmental conditions known for their effects on gene expression related to Δ (see above). QTL analysis was repeated on several completely
 10 independent data sets obtained under highly controlled conditions in glasshouses or growth chambers, where either air humidity, photoperiod or irradiance (amount, diurnal pattern, day to day variation) was varied. Depending on the experiment, all 100 recombinants inbred lines were included or only the subset of lines with cross-overs on chromosome 2. These experiments confirmed that genetic variation in Δ
 15 could be mostly ascribed to a portion of chromosome 2 (Table 1) between about 46-50.7 cM.

- When RILs were sorted graphically according to carbon isotope discrimination and their genotype at the *ER* marker (50.64 cM) and its vicinity (*Ld-er1* genotype or Col-
 20 *ER* genotype), lines which were *Ld-er* at the *ERECTA* marker ranked mostly at the high end of carbon isotope discrimination values. In contrast, lines having a Col-*ERECTA* marker genotype ranked mostly at the low end of carbon isotope discrimination values (data available on request). In the middle of the range of carbon isotope discrimination values, there was some overlap between the two sets
 25 of lines. Some lines were always at an extreme (in all 18 experiments performed), while the ranking of other lines was more unstable. These data indicate a locus for transpiration efficiency, as determined by the carbon isotope discrimination value, in the vicinity of the *ERECTA* locus on chromosome 2 (Table 1). This locus most likely involves the *ER* gene. Depending on the positions of cross-overs between *Ld-er*
 30 and *Col*, recombination between *ERECTA* and one or more of the other genes influences the transpiration efficiency phenotype of the progeny.

EXAMPLE 5

Determination of a role for the *ERECTA* gene in regulating transpiration efficiency

We compared Col and Ler ecotypes with near-isogenic mutant lines for the *erecta* gene, to examine a possible role of the *ERECTA* gene in determining carbon isotope discrimination (Δ).

- 5 Plants expressing the wild type *ERECTA* gene (SEQ ID NO: 1), or an *erecta* mutant allele in the Columbia background (eg. Col-*er101* to -*er105*; or Col-*er108* to -*er123*), have been publicly described. Two of these mutants were available for comparison to the isogenic or near-isogenic lines (Table 2).
- 10 Col4, the other parental line (*ER*) for Lister and Dean's RILs was systematically included in the comparison. Where possible, other Col "ecotypes" were also included, (eg. Col0, Col3-7), to assess their similarity with respect to carbon isotope discrimination, especially compared to the RIL parental ecotype Col4.
- 15 The results of these comparisons are described in Table 3. Data indicate that the differences in carbon isotope discrimination values between *er* and *ER* lines for 15 different experimental runs corresponding to growth under low to high light (100 to 800 $\mu\text{E m}^{-2} \text{ s}^{-1}$), low to high humidity (40 to 85%), short to long days (8, 10, 24hrs), normal to high temperatures (22/20°C to 28/20 °C).
- 20 As expected, the spread of carbon isotope discrimination values among lines varied with environmental conditions. Lines carrying *er* mutations have a greater carbon isotope discrimination value overall than those having the *ER* wild type gene (see Table 3, column 1), indicative of a lower water use-efficiency. There is usually little
- 25 difference in C isotopic discrimination between the various Col lines, (see the similar averages obtained for columns 2, 3, and 4 in Table 3, wherein *er105* is compared to 3 different Col ecotypes, Col0, Col4 and 3176 or Col1). When present, the *er105* mutant always has the greatest carbon isotope discrimination value of all lines, including *er1* and *er2* (columns 2-4 compared to columns 5-6 in Table 3, or
- 30 column 8 compared to column 9 in Table 3). The value measured in the *er105* mutant is always significantly greater than in the *ER* isogenic line (column 4 in Table 3). The value measured in *er1* (Landsberg parental line NW20) is usually also greater than that in the *ER* lines 3177 (near isogenic, column 6 of Table 3), and to a lesser extent Col4 (Columbia parental line, column 7 of Table 3). These observations give

direct evidence that the *ERECTA* gene plays a significant role in determining genetic differences in carbon isotopic discrimination in *Arabidopsis*.

This conclusion is independently confirmed by leaf gas exchange measurements that allow the direct measure of transpiration efficiency (ratio of net CO₂ fixation to water loss; column 4 in Table 4; Figures 1a-1c, 2a-2c). Measurements on mature leaves reveal that *ER* lines are characterised by a greater ratio of CO₂ assimilation to water loss than lines carrying *er* mutations. This is most obvious when comparing the pair *Col1/er105* with a 21% greater transpiration efficiency (ratio A/E) in *Col1* than *er105*, or the pair *Col1/er2* with a 16% greater transpiration efficiency in *Col1*. Consistent with the measurements of carbon isotope discrimination, the effect *er/ER* is relatively smaller in the *Ld* background (9% greater ratio A/E in *Ld-ER* (3177) than the *Ld-er1*(NSW20) background.

Also consistent with the carbon discrimination measurements, is the 20% difference in transpiration efficiency between the two RILs parental lines (4.06 and 3.38 mmolC/molH₂O in *Col4-ER* and *Ld-er1*, respectively).

The fact that of all 3 *erecta* mutants examined, *er105* has the most extreme carbon discrimination and transpiration efficiency phenotypes suggests that the *er105* mutation affects a more crucial part of the *ERECTA* gene than *er2* or *er1*. This is consistent with the published data on the *er105* mutant. This mutation corresponds to the insertion of a large "foreign insert" in the *ERECTA* gene (1200bp). The insertion totally inhibits transcription of the gene and causes the strongest *erecta* phenotype of all *erecta* mutants isolated in *Col* (with respect to inflorescence clustering and silique width and shape. Alternatively, or in addition, data indicate that *erecta* mutations have a stronger effect on carbon isotope discrimination values in a Columbia genetic background than in a Landsberg background (comparison of phenotypic effects of *er105* and *er1*), implying that other genes, polymorphic between Landsberg and Columbia ecotypes, interact with *ERECTA* in determining transpiration efficiency. This could also account for the greater difference in transpiration efficiency between *er/ER* lines in *Col* background than in a *Ld* background (see above, Table 4). Alternatively, or in addition, data indicate that the *erecta* mutation is not the only mutation present in the *er105* mutant. For example,

the mutagenized Col seeds may have carried the *gl1* mutation, induced by the fast neutron irradiation, that also contributes to the phenotype observed.

A comparison of transcript profiles in *er/ER* isogenic lines (in both Col and Ld
5 background) allows determination of the involvement of additional genes to *ERECTA* and the effect of environment on their expression.

TABLE 1

QTL Analysis of Carbon Isotope Discrimination in Lister and Dean's Recombinant Inbred Lines

RUN No.	chr2 locus	QTL analysis method	chr4 locus	QTLs number	CONCLUSION predicted map position
Experimental conditions					
Run 1 (40 lines) Glasshouse-					
12h day length	58.5	SIM&CIM		2	chr2: 58.5-61.02
irradiance 150-350 $\mu\text{E m}^{-2} \text{s}^{-1}$	46.77	SIM&CIM			chr2: 46.77-50.75
Seedlings transferred from agar plates	61.02	SIM	108.5		
Run1 data but with using different markers					
	56.94 to 58.00	CIM&SIM		1	
	46.77 to 50.75	SIM			
	63.02				

RUN No.	chr2 locus	QTL analysis method	chr4 locus	QTLs number	CONCLUSION predicted map position
Experimental conditions					
Run 1					
with different number of lines	58.5 to 61.02			2	
	56-61				
Run2					
Glasshouse					
September					
from seeds sown on soil					
batch 1	50.75	CIM (QTL cart)		2	chr2: 56.94-61.02
	61.02	MQTL			chr2: 50.75
batch 2	?50.75	MQTL		NS	
batch3-5					
all batches	58.5	MQTLcart		NS	
	56.94-58.5	MQTL			

RUN No.	chr2 locus	QTL analysis method	chr4 locus	CONCLUSION
Experimental conditions				QTLs number predicted map position
Run 3				
37 lines: parents and lines with crossing-overs on chromosome 2				
5 growth conditions differing in humidity, irradiance, mode of establishment (seeds sown on soil or seedlings transplanted from agar)				
batch B	61.02-61.06		108 NS	
batch C	56.94-58.00			
batch D	63.02	QTLcar		
	63.02	MQTL		
all batches (conditions)	58.5			1 or 2? chr2: 56.94-58.5
	61.02			3 chr2: 61.02-63.02

RUN No.	chr2 locus	QTL analysis method	chr4 locus	QTLs number	CONCLUSION predicted map position
Run 4					
same lines as Run 3					
growth chambers					
10h daylight					
	50.74				chr2: 50.74
Run 5					
repeat of run 1 BUT ALL lines					
	50.74			1	chr2: 50.74
Run 7					
same lines as in run 1 but in growth chamber and higher light					
10h daylight					
470-510 $\mu\text{E m}^{-2} \text{s}^{-1}$ irradiance					
	46.77-50.75	CIM&SIM		1	chr 2: 46.77-5065

TABLE 2

Background	Mutation	Stock Centre name	Isogenic <i>ER</i> line and Stock Centre Name
Landsberg	<i>er1</i>	CS20 or NW20 ^a	3177 or CS163
Columbia	<i>er2</i> ^b	3401	Col1 or 3176
Columbia	<i>er105</i> ^c		Col3 with gl1 marker or Col0

- a, NW20 is an *Ler* parent for Lister and Dean's recombinant lines, carrying the *er1* mutation. Lines 3177 or CS163 are the closest isogenic ER lines.
- b, *er2* is an *er* allele identified by R       in Col background. Col1 or 3176 are the closest Col near-isogenic lines.
- c, *er105* was isolated from a fast-neutron-irradiated Col seed population (Torii *et al.*, 1996).
- 10 d, Col4, the *Col* parent for the Lister and Dean's parent was systemically included in all comparisons.

TABLE 3

Comparison of er/ER lines in both Col and Ld background for carbon isotope discrimination values
(per ml) in leaf material under a range of environmental conditions

Run No.	Differences in mean carbon isotope discrimination values (per mil)								
	(1) er-ER (all lines)	(2) er105-Col0	(3) er105-Col4	(4) er105-3176	(5) er2-3176	(6) er1-3177	(7) er1-Col4 (parental lines for RLs)	(8) er105-Coli	(9) er1-Coli
1	0.13	0.16						0.16	
2	0.89	1.18						1.18	
3	0.26	0.11						0.11	
4	1.12	1.60						1.60	
5	1.03		1.83	1.67	0.92	0.64	0.82	1.75	0.73
6	0.70	1.13	1.01	0.71	0.27	0.74	0.73	0.95	0.73
7	0.70	1.32	1.12	1.23	0.75	0.35	0.05	1.22	0.15
8	0.59	1.16	1.11	1.19	0.54	0.28	0.06	1.16	0.17
9	0.30	1.09	0.77	0.77	0.02	0.00	0.56	0.88	0.28
10	0.56	1.05	0.94	0.87	0.38	0.39	0.33	0.95	0.36

11	0.48			0.40			0.52	0.40	0.52
12	0.36	0.82		1.31	1.08	0.33	0.05	1.07	0.01
13	0.38			0.90	0.82	0.07	0.60	0.86	0.56
14	0.65	1.42		0.60			0.58	1.01	0.32
15	0.82								0.82

For all runs:

Mean	0.60	1.01	1.00	1.04	0.41	0.40	0.41	0.95	0.42
S.E.	0.07	0.14	0.11	0.11	0.10	0.08	0.09	0.12	0.08

For Common runs:

Mean:	0.58	1.10	1.12	1.04	0.41	0.38	0.39	1.11	0.37
S.E.	0.08	0.05	0.11	0.11	0.10	0.09	0.10	0.10	0.09

TABLE 4

Run 9- December 2001: Leaf gas exchange measurements in er/ER Arabidopsis lines

Genotype	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)		
	E	A	Gw	A/E	pa	pi	pi/pa	1-pi/pa		
	mmol H ₂ O /m ² /s		mol/m ² /s	mmolC/ μbar	μbar					
Row (1) Ld-ER	3177-G	Mean	3.38	12.33	0.273	3.67	360	282	0.782	0.218
		S.E.	0.48	1.64	0.039	0.14	10	11	0.010	0.010
Row (2) Ld-er	NSW20 E	Mean	2.59	8.73	0.218	3.38	348	280	0.804	0.196
		S.E.	0.07	0.31	0.005	0.04	5	4	0.002	0.002
Row (3) Col-ER	933	Mean	3.41	13.55	0.291	4.06	350	270	0.772	0.228
		S.E.	0.40	1.16	0.040	0.22	4	7	0.020	0.020
Row (4) Col-ER	3176 J	Mean	2.23	10.13	0.180	4.55	346	254	0.734	0.266
	(Col1)	S.E.	0.50	1.47	0.048	0.24	5	9	0.021	0.021

Row (5) Col-er	er105 F	Mean	2.27	8.55	0.198	3.76	356	283	0.795	0.205
		S.E.	0.03	0.17	0.005	0.07	11	10	0.006	0.006
Row (6) Col-er	er2C	Mean	3.06	11.90	0.256	3.92	357	279	0.780	0.220
		S.E.	0.22	0.56	0.027	0.12	1	6	0.014	0.014

CONCLUSION

Comparison Ld-ER/Ld-er	er line has lower A/E with lower g and lower A The difference in A/E is driven by A
Comparison 933/NSW20	NSW20 (er) has lower A/E with lower g and lower A The difference in A/E is driven by A
Comparison Col1/Ld-er1	er105 has MUCH lower A/E with Higher g and lower A i.e. the difference in A/E is driven by A and g
Comparison Col1/Col-er105	er2 has lower A/E with MUCH higher g and HIGHER A i.e. the difference in A/E is driven by g and is opposed or not driven by A

NOTE: p_a and p_l are the ambient and intercellular partial pressures of CO_2 , respectively.

SEQUENCE LISTING

5
 <110> The Australian National University
 Masle, Josette
10 Farquhar, Graham
 Gilmore, Scott

15
 <120> Method of producing plants having enhanced transpiration
 efficiency and plants produced therefrom I

20
 <130> 500592/MRO

25 <160> 10

30 <170> PatentIn version 3.1

 <210> 1
35 <211> 3176
 <212> DNA
40 <213> Arabidopsis thaliana ERECTA allele

 <220>
45 <221> CDS
 <222> (77)..(3004)
 <223>
50

 <400> 1
55 gtttcttctt catggagact tgaaagcttt taaagtatat ctaaaaacgc agtcgtttta
60

agactgtgtg tgagaa atg gct ctg ttt aga gat att gtt ctt ctt ggg ttt
 112
 Met Ala Leu Phe Arg Asp Ile Val Leu Leu Gly Phe
 1 5 10
 5
 ctc ttc tgc ttg agc tta gta gct act gtg act tca gag gag gga gca
 160
 Leu Phe Cys Leu Ser Leu Val Ala Thr Val Thr Ser Glu Glu Gly Ala
 15 20 25
 10
 acg ttg ctg gag att aag aag tca ttc aaa gat gtg aac aat gtt ctt
 208
 Thr Leu Leu Glu Ile Lys Lys Ser Phe Lys Asp Val Asn Asn Val Leu
 30 35 40
 15
 tat gac tgg aca act tca cct tct tcg gat tat tgt gtc tgg aga ggt
 256
 Tyr Asp Trp Thr Thr Ser Pro Ser Ser Asp Tyr Cys Val Trp Arg Gly
 45 50 55 60
 20
 gtg tct tgt gaa aat gtc acc ttc aat gtt gtt gct ctt aat ttg tca
 304
 Val Ser Cys Glu Asn Val Thr Phe Asn Val Val Ala Leu Asn Leu Ser
 65 70 75
 25
 gat ttg aat ctt gat gga gaa atc tca cct gct att gga gat ctc aag
 352
 Asp Leu Asn Leu Asp Gly Glu Ile Ser Pro Ala Ile Gly Asp Leu Lys
 80 85 90
 30
 agt ctc ttg tca att gat ctg cga ggt aat cgc ttg tct gga caa atc
 400
 Ser Leu Leu Ser Ile Asp Leu Arg Gly Asn Arg Leu Ser Gly Gln Ile
 95 100 105
 35
 cct gat gag att ggt gac tgt tct tct ttg caa aac tta gac tta tcc
 448
 Pro Asp Glu Ile Gly Asp Cys Ser Ser Leu Gln Asn Leu Asp Leu Ser
 110 115 120
 40
 ttc aat gaa tta agt ggt gac ata ccg ttt tcg att tcg aag ttg aag
 496
 Phe Asn Glu Leu Ser Gly Asp Ile Pro Phe Ser Ile Ser Lys Leu Lys
 125 130 135 140
 45
 caa ctt gag cag ctg att ctg aag aat aac caa ttg ata gga ccg atc
 544
 Gln Leu Glu Gln Leu Ile Leu Lys Asn Asn Gln Leu Ile Gly Pro Ile
 145 150 155
 50
 cct tca aca ctt tca cag att cca aac ctg aaa att ctg gac ttg gca
 592
 Pro Ser Thr Leu Ser Gln Ile Pro Asn Leu Lys Ile Leu Asp Leu Ala
 160 165 170
 55
 cag aat aaa ctc agt ggt gag ata cca aga ctt att tac tgg aat gaa
 640

Gln Asn Lys Leu Ser Gly Glu Ile Pro Arg Leu Ile Tyr Trp Asn Glu
 175 180 185
 5 gtt ctt cag tat ctt ggg ttg cga gga aac aac tta gtc ggt aac att
 688
 Val Leu Gln Tyr Leu Gly Leu Arg Gly Asn Asn Leu Val Gly Asn Ile
 190 195 200
 10 tct cca gat ttg tgt caa ctg act ggt ctt tgg tat ttt gac gta aga
 736
 Ser Pro Asp Leu Cys Gln Leu Thr Gly Leu Trp Tyr Phe Asp Val Arg
 205 210 215 220
 15 aac aac agt ttg act ggt agt ata cct gag acg ata gga aat tgc act
 784
 Asn Asn Ser Leu Thr Gly Ser Ile Pro Glu Thr Ile Gly Asn Cys Thr
 225 230 235
 20 gcc ttc cag gtt ttg gac ttg tcc tac aat cag cta act ggt gag atc
 832
 Ala Phe Gln Val Leu Asp Leu Ser Tyr Asn Gln Leu Thr Gly Glu Ile
 240 245 250
 25 cct ttt gac atc ggc ttc ctg caa gtt gca aca tta tca ttg caa ggc
 880
 Pro Phe Asp Ile Gly Phe Leu Gln Val Ala Thr Leu Ser Leu Gln Gly
 255 260 265
 30 aat caa ctc tct ggg aag att cca tca gtg att ggt ctc atg caa gcc
 928
 Asn Gln Leu Ser Gly Lys Ile Pro Ser Val Ile Gly Leu Met Gln Ala
 270 275 280
 35 ctt gca gtc tta gat cta agt ggc aac ttg ttg agt gga tct att cct
 976
 Leu Ala Val Leu Asp Leu Ser Gly Asn Leu Leu Ser Gly Ser Ile Pro
 285 290 295 300
 40 ccg att ctc gga aat ctt act ttc acc gag aaa ttg tat ttg cac agt
 1024
 Pro Ile Leu Gly Asn Leu Thr Phe Thr Glu Lys Leu Tyr Leu His Ser
 305 310 315
 45 aac aag ctg act ggt tca att cca cct gag ctt gga aac atg tca aaa
 1072
 Asn Lys Leu Thr Gly Ser Ile Pro Pro Glu Leu Gly Asn Met Ser Lys
 320 325 330
 50 ctc cat tac ctg gaa ctc aat gat aat cat ctc acg ggt cat ata cca
 1120
 Leu His Tyr Leu Glu Leu Asn Asp Asn His Leu Thr Gly His Ile Pro
 335 340 345
 55 cca gag ctt ggg aag ctt act gac ttg ttt gat ctg aat gtg gcc aac
 1168
 Pro Glu Leu Gly Lys Leu Thr Asp Leu Phe Asp Leu Asn Val Ala Asn
 350 355 360

aat gat ctg gaa gga cct ata cct gat cat ctg agc tct tgc aca aat
 1216
 Asn Asp Leu Glu Gly Pro Ile Pro Asp His Leu Ser Ser Cys Thr Asn
 5 365 370 375 380

 cta aac agc tta aat gtt cat ggg aac aag ttt agt ggc act ata ccc
 1264
 Leu Asn Ser Leu Asn Val His Gly Asn Lys Phe Ser Gly Thr Ile Pro
 10 385 390 395

 cga gca ttt caa aag cta gaa agt atg act tac ctt aat ctg tcc agc
 1312
 Arg Ala Phe Gln Lys Leu Glu Ser Met Thr Tyr Leu Asn Leu Ser Ser
 15 400 405 410

 aac aat atc aaa ggt cca atc ccg gtt gag cta tct cgt atc ggt aac
 1360
 Asn Asn Ile Lys Gly Pro Ile Pro Val Glu Leu Ser Arg Ile Gly Asn
 20 415 420 425

 tta gat aca ttg gat ctt tcc aac aac aag ata aat gga atc att cct
 1408
 Leu Asp Thr Leu Asp Leu Ser Asn Asn Lys Ile Asn Gly Ile Ile Pro
 25 430 435 440

 tct tcc ctt ggt gat ttg gag cat ctt ctc aag atg aac ttg agt aga
 1456
 Ser Ser Leu Gly Asp Leu Glu His Leu Leu Lys Met Asn Leu Ser Arg
 30 445 450 455 460

 aat cat ata act ggt gta gtt cca ggc gac ttt gga aat cta aga agc
 1504
 Asn His Ile Thr Gly Val Val Pro Gly Asp Phe Gly Asn Leu Arg Ser
 35 465 470 475

 atc atg gaa ata gat ctt tca aat aat gat atc tct ggc cca att cca
 1552
 Ile Met Glu Ile Asp Leu Ser Asn Asn Asp Ile Ser Gly Pro Ile Pro
 40 480 485 490

 gaa gag ctt aac caa tta cag aac ata att ttg ctg aga ctg gaa aat
 1600
 Glu Glu Leu Asn Gln Leu Gln Asn Ile Ile Leu Leu Arg Leu Glu Asn
 45 495 500 505

 aat aac ctg act ggt aat gtt ggt tca tta gcc aac tgt ctc agt ctc
 1648
 Asn Asn Leu Thr Gly Asn Val Gly Ser Leu Ala Asn Cys Leu Ser Leu
 50 510 515 520

 act gta ttg aat gta tct cat aac aac ctc gta ggt gat atc cct aag
 1696
 Thr Val Leu Asn Val Ser His Asn Asn Leu Val Gly Asp Ile Pro Lys
 55 525 530 535 540

aac aat aac ttc tca aga ttt tca cca gac agc ttc att ggc aat cct
 1744
 Asn Asn Asn Phe Ser Arg Phe Ser Pro Asp Ser Phe Ile Gly Asn Pro
 545 550 555
 5
 ggt ctt tgc ggt agt tgg cta aac tca ccg tgt cat gat tct cgt cga
 1792
 Gly Leu Cys Gly Ser Trp Leu Asn Ser Pro Cys His Asp Ser Arg Arg
 560 565 570
 10
 act gta cga gtg tca atc tct aga gca gct att ctt gga ata gct att
 1840
 Thr Val Arg Val Ser Ile Ser Arg Ala Ala Ile Leu Gly Ile Ala Ile
 575 580 585
 15
 ggg gga ctt gtg atc ctt ctc atg gtc tta ata gca gct tgc cga ccg
 1888
 Gly Gly Leu Val Ile Leu Leu Met Val Leu Ile Ala Ala Cys Arg Pro
 590 595 600
 20
 cat aat cct cct cct ttt ctt gat gga tca ctt gac aaa cca gta act
 1936
 His Asn Pro Pro Pro Phe Leu Asp Gly Ser Leu Asp Lys Pro Val Thr
 605 610 615 620
 25
 tat tcg aca ccg aag ctc gtc atc ctt cat atg aac atg gca ctc cac
 1984
 Tyr Ser Thr Pro Lys Leu Val Ile Leu His Met Asn Met Ala Leu His
 625 630 635
 30
 gtt tac gag gat atc atg aga atg aca gag aat cta agt gag aag tat
 2032
 Val Tyr Glu Asp Ile Met Arg Met Thr Glu Asn Leu Ser Glu Lys Tyr
 640 645 650
 35
 atc att ggg cac gga gca tca agc act gta tac aaa tgt gtt ttg aag
 2080
 Ile Ile Gly His Gly Ala Ser Ser Thr Val Tyr Lys Cys Val Leu Lys
 655 660 665
 40
 aat tgt aaa ccg gtt gcg att aag cgg ctt tac tct cac aac cca cag
 2128
 Asn Cys Lys Pro Val Ala Ile Lys Arg Leu Tyr Ser His Asn Pro Gln
 670 675 680
 45
 tca atg aaa cag ttt gaa aca gaa ctc gag atg cta agt agc atc aag
 2176
 Ser Met Lys Gln Phe Glu Thr Glu Leu Glu Met Leu Ser Ser Ile Lys
 685 690 695 700
 50
 cac aga aat ctt gtg agc cta caa gct tat tcc ctc tct cac ttg ggg
 2224
 His Arg Asn Leu Val Ser Leu Gln Ala Tyr Ser Leu Ser His Leu Gly
 705 710 715
 55
 agt ctt ctg ttc tat gac tat ttg gaa aat ggt agc ctc tgg gat ctt
 2272

Ser Leu Leu Phe Tyr Asp Tyr Leu Glu Asn Gly Ser Leu Trp Asp Leu
 720 725 730
 5 ctt cat ggc cct acg aag aaa aag act ctt gat tgg gac aca cgg ctt
 2320
 Leu His Gly Pro Thr Lys Lys Lys Thr Leu Asp Trp Asp Thr Arg Leu
 735 740 745
 10 aag ata gca tat ggt gca gca caa ggt tta gct tat cta cac cat gac
 2368
 Lys Ile Ala Tyr Gly Ala Ala Gln Gly Leu Ala Tyr Leu His His Asp
 750 755 760
 15 tgt agt cca agg atc att cac aga gac gtg aag tcg tcc aac att ctc
 2416
 Cys Ser Pro Arg Ile Ile His Arg Asp Val Lys Ser Ser Asn Ile Leu
 765 770 775 780
 20 ttg gac aaa gac tta gag gct cgt ttg aca gat ttt gga ata gcg aaa
 2464
 Leu Asp Lys Asp Leu Glu Ala Arg Leu Thr Asp Phe Gly Ile Ala Lys
 785 790 795
 25 agc ttg tgt gtg tca aag tca cat act tca act tac gtg atg ggc acg
 2512
 Ser Leu Cys Val Ser Lys Ser His Thr Ser Thr Tyr Val Met Gly Thr
 800 805 810
 30 ata ggt tac ata gac ccc gag tat gct cgc act tca cgg ctc act gag
 2560
 Ile Gly Tyr Ile Asp Pro Glu Tyr Ala Arg Thr Ser Arg Leu Thr Glu
 815 820 825
 35 aaa tcc gat gtc tac agt tat gga ata gtc ctt ctt gag ttg tta acc
 2608
 Lys Ser Asp Val Tyr Ser Tyr Gly Ile Val Leu Leu Glu Leu Leu Thr
 830 835 840
 40 cga agg aaa gcc gtt gat gac gaa tcc aat ctc cac cat ctg ata atg
 2656
 Arg Arg Lys Ala Val Asp Asp Glu Ser Asn Leu His His Leu Ile Met
 845 850 855 860
 45 tca aag acg ggg aac aat gaa gtg atg gaa atg gca gat cca gac atc
 2704
 Ser Lys Thr Gly Asn Asn Glu Val Met Glu Met Ala Asp Pro Asp Ile
 865 870 875
 50 aca tcg acg tgt aaa gat ctc ggt gtg gtg aag aaa gtt ttc caa ctg
 2752
 Thr Ser Thr Cys Lys Asp Leu Gly Val Val Lys Lys Val Phe Gln Leu
 880 885 890
 55 gca ctc cta tgc acc aaa aga cag ccg aat gat cga ccc aca atg cac
 2800
 Ala Leu Leu Cys Thr Lys Arg Gln Pro Asn Asp Arg Pro Thr Met His
 895 900 905

cag gtg act cgt gtt ctc ggc agt ttt atg cta tcg gaa caa cca cct
 2848
 5 Gln Val Thr Arg Val Leu Gly Ser Phe Met Leu Ser Glu Gln Pro Pro
 910 915 920

gct gcg act gac acg tca gcg acg ctg gct ggt tcg tgc tac gtc gat
 2896
 10 Ala Ala Thr Asp Thr Ser Ala Thr Leu Ala Gly Ser Cys Tyr Val Asp
 925 930 935 940

gag tat gca aat ctc aag act cct cat tct gtc aat tgc tct tcc atg
 2944
 15 Glu Tyr Ala Asn Leu Lys Thr Pro His Ser Val Asn Cys Ser Ser Met
 945 950 955

agt gct tct gat gct caa ctg ttt ctt cgg ttt gga caa gtt att tct
 2992
 20 Ser Ala Ser Asp Ala Gln Leu Phe Leu Arg Phe Gly Gln Val Ile Ser
 960 965 970

cag aac agt gag tagtttttcg ttaggaggag aatctttaaa acggtatctt
 3044
 25 Gln Asn Ser Glu
 975

ttcggtgcgt taagctgtta gaaaaattaa tgtctcatgt aaagtattat gcactgcctt
 3104

30 attattatta gacaagtgtg tgggtgtgaat atgtcttcag actggcactt agacttccta
 3164

taagttcttg cc
 3176
 35

<210> 2

<211> 976
 40

<212> PRT

<213> Arabidopsis thaliana ERECTA allele

45

<400> 2

50 Met Ala Leu Phe Arg Asp Ile Val Leu Leu Gly Phe Leu Phe Cys Leu
 1 5 10 15

Ser Leu Val Ala Thr Val Thr Ser Glu Glu Gly Ala Thr Leu Leu Glu
 20 25 30
 55

Ile Lys Lys Ser Phe Lys Asp Val Asn Asn Val Leu Tyr Asp Trp Thr

	35		40		45												
5	Thr	Ser	Pro	Ser	Ser	Asp	Tyr	Cys	Val	Trp	Arg	Gly	Val	Ser	Cys	Glu	
	50						55					60					
10	Asn	Val	Thr	Phe	Asn	Val	Val	Ala	Leu	Asn	Leu	Ser	Asp	Leu	Asn	Leu	
	65					70					75					80	
15	Asp	Gly	Glu	Ile	Ser	Pro	Ala	Ile	Gly	Asp	Leu	Lys	Ser	Leu	Leu	Ser	
					85					90					95		
20	Ile	Asp	Leu	Arg	Gly	Asn	Arg	Leu	Ser	Gly	Gln	Ile	Pro	Asp	Glu	Ile	
				100					105					110			
25	Gly	Asp	Cys	Ser	Ser	Leu	Gln	Asn	Leu	Asp	Leu	Ser	Phe	Asn	Glu	Leu	
			115					120					125				
30	Ser	Gly	Asp	Ile	Pro	Phe	Ser	Ile	Ser	Lys	Leu	Lys	Gln	Leu	Glu	Gln	
	130						135					140					
35	Leu	Ile	Leu	Lys	Asn	Asn	Gln	Leu	Ile	Gly	Pro	Ile	Pro	Ser	Thr	Leu	
	145					150					155					160	
40	Ser	Gln	Ile	Pro	Asn	Leu	Lys	Ile	Leu	Asp	Leu	Ala	Gln	Asn	Lys	Leu	
					165					170					175		
45	Ser	Gly	Glu	Ile	Pro	Arg	Leu	Ile	Tyr	Trp	Asn	Glu	Val	Leu	Gln	Tyr	
				180					185					190			
50	Leu	Gly	Leu	Arg	Gly	Asn	Asn	Leu	Val	Gly	Asn	Ile	Ser	Pro	Asp	Leu	
			195					200					205				
55	Cys	Gln	Leu	Thr	Gly	Leu	Trp	Tyr	Phe	Asp	Val	Arg	Asn	Asn	Ser	Leu	
	210						215					220					
60	Thr	Gly	Ser	Ile	Pro	Glu	Thr	Ile	Gly	Asn	Cys	Thr	Ala	Phe	Gln	Val	
	225					230					235					240	
65	Leu	Asp	Leu	Ser	Tyr	Asn	Gln	Leu	Thr	Gly	Glu	Ile	Pro	Phe	Asp	Ile	
					245					250					255		
70	Gly	Phe	Leu	Gln	Val	Ala	Thr	Leu	Ser	Leu	Gln	Gly	Asn	Gln	Leu	Ser	
				260					265					270			

Gly Lys Ile Pro Ser Val Ile Gly Leu Met Gln Ala Leu Ala Val Leu
 275 280 285
 5

Asp Leu Ser Gly Asn Leu Leu Ser Gly Ser Ile Pro Pro Ile Leu Gly
 290 295 300

10

Asn Leu Thr Phe Thr Glu Lys Leu Tyr Leu His Ser Asn Lys Leu Thr
 305 310 315 320

15

Gly Ser Ile Pro Pro Glu Leu Gly Asn Met Ser Lys Leu His Tyr Leu
 325 330 335

20

Glu Leu Asn Asp Asn His Leu Thr Gly His Ile Pro Pro Glu Leu Gly
 340 345 350

Lys Leu Thr Asp Leu Phe Asp Leu Asn Val Ala Asn Asn Asp Leu Glu
 355 360 365

25

Gly Pro Ile Pro Asp His Leu Ser Ser Cys Thr Asn Leu Asn Ser Leu
 370 375 380

30

Asn Val His Gly Asn Lys Phe Ser Gly Thr Ile Pro Arg Ala Phe Gln
 385 390 395 400

35

Lys Leu Glu Ser Met Thr Tyr Leu Asn Leu Ser Ser Asn Asn Ile Lys
 405 410 415

40

Gly Pro Ile Pro Val Glu Leu Ser Arg Ile Gly Asn Leu Asp Thr Leu
 420 425 430

45

Asp Leu Ser Asn Asn Lys Ile Asn Gly Ile Ile Pro Ser Ser Leu Gly
 435 440 445

46

Asp Leu Glu His Leu Leu Lys Met Asn Leu Ser Arg Asn His Ile Thr
 450 455 460

50

Gly Val Val Pro Gly Asp Phe Gly Asn Leu Arg Ser Ile Met Glu Ile
 465 470 475 480

55

Asp Leu Ser Asn Asn Asp Ile Ser Gly Pro Ile Pro Glu Glu Leu Asn
 485 490 495

	Gln	Leu	Gln	Asn	Ile	Ile	Leu	Leu	Arg	Leu	Glu	Asn	Asn	Asn	Leu	Thr	
				500					505						510		
5	Gly	Asn	Val	Gly	Ser	Leu	Ala	Asn	Cys	Leu	Ser	Leu	Thr	Val	Leu	Asn	
			515					520					525				
10	Val	Ser	His	Asn	Asn	Leu	Val	Gly	Asp	Ile	Pro	Lys	Asn	Asn	Asn	Phe	
		530					535					540					
15	Ser	Arg	Phe	Ser	Pro	Asp	Ser	Phe	Ile	Gly	Asn	Pro	Gly	Leu	Cys	Gly	
	545					550					555					560	
20	Ser	Trp	Leu	Asn	Ser	Pro	Cys	His	Asp	Ser	Arg	Arg	Thr	Val	Arg	Val	
				565						570					575		
25	Ser	Ile	Ser	Arg	Ala	Ala	Ile	Leu	Gly	Ile	Ala	Ile	Gly	Gly	Leu	Val	
			580						585					590			
30	Ile	Leu	Leu	Met	Val	Leu	Ile	Ala	Ala	Cys	Arg	Pro	His	Asn	Pro	Pro	
		595						600					605				
35	Pro	Phe	Leu	Asp	Gly	Ser	Leu	Asp	Lys	Pro	Val	Thr	Tyr	Ser	Thr	Pro	
		610					615					620					
40	Lys	Leu	Val	Ile	Leu	His	Met	Asn	Met	Ala	Leu	His	Val	Tyr	Glu	Asp	
	625					630					635					640	
45	Ile	Met	Arg	Met	Thr	Glu	Asn	Leu	Ser	Glu	Lys	Tyr	Ile	Ile	Gly	His	
				645						650					655		
50	Gly	Ala	Ser	Ser	Thr	Val	Tyr	Lys	Cys	Val	Leu	Lys	Asn	Cys	Lys	Pro	
			660						665					670			
55	Val	Ala	Ile	Lys	Arg	Leu	Tyr	Ser	His	Asn	Pro	Gln	Ser	Met	Lys	Gln	
		675						680						685			
60	Phe	Glu	Thr	Glu	Leu	Glu	Met	Leu	Ser	Ser	Ile	Lys	His	Arg	Asn	Leu	
		690					695					700					
65	Val	Ser	Leu	Gln	Ala	Tyr	Ser	Leu	Ser	His	Leu	Gly	Ser	Leu	Leu	Phe	
	705					710					715					720	

	Tyr	Asp	Tyr	Leu	Glu	Asn	Gly	Ser	Leu	Trp	Asp	Leu	Leu	His	Gly	Pro	
					725					730					735		
5	Thr	Lys	Lys	Lys	Thr	Leu	Asp	Trp	Asp	Thr	Arg	Leu	Lys	Ile	Ala	Tyr	
				740					745					750			
10	Gly	Ala	Ala	Gln	Gly	Leu	Ala	Tyr	Leu	His	His	Asp	Cys	Ser	Pro	Arg	
			755					760					765				
15	Ile	Ile	His	Arg	Asp	Val	Lys	Ser	Ser	Asn	Ile	Leu	Leu	Asp	Lys	Asp	
		770					775					780					
20	Leu	Glu	Ala	Arg	Leu	Thr	Asp	Phe	Gly	Ile	Ala	Lys	Ser	Leu	Cys	Val	
	785					790					795					800	
25	Ser	Lys	Ser	His	Thr	Ser	Thr	Tyr	Val	Met	Gly	Thr	Ile	Gly	Tyr	Ile	
					805					810					815		
30	Asp	Pro	Glu	Tyr	Ala	Arg	Thr	Ser	Arg	Leu	Thr	Glu	Lys	Ser	Asp	Val	
				820					825					830			
35	Tyr	Ser	Tyr	Gly	Ile	Val	Leu	Leu	Glu	Leu	Leu	Thr	Arg	Arg	Lys	Ala	
			835				840						845				
40	Val	Asp	Asp	Glu	Ser	Asn	Leu	His	His	Leu	Ile	Met	Ser	Lys	Thr	Gly	
	850						855					860					
45	Asn	Asn	Glu	Val	Met	Glu	Met	Ala	Asp	Pro	Asp	Ile	Thr	Ser	Thr	Cys	
	865					870					875					880	
50	Lys	Asp	Leu	Gly	Val	Val	Lys	Lys	Val	Phe	Gln	Leu	Ala	Leu	Leu	Cys	
					885					890					895		
55	Thr	Lys	Arg	Gln	Pro	Asn	Asp	Arg	Pro	Thr	Met	His	Gln	Val	Thr	Arg	
				900					905					910			
60	Val	Leu	Gly	Ser	Phe	Met	Leu	Ser	Glu	Gln	Pro	Pro	Ala	Ala	Thr	Asp	
		915					920						925				
65	Thr	Ser	Ala	Thr	Leu	Ala	Gly	Ser	Cys	Tyr	Val	Asp	Glu	Tyr	Ala	Asn	
	930						935					940					
70	Leu	Lys	Thr	Pro	His	Ser	Val	Asn	Cys	Ser	Ser	Met	Ser	Ala	Ser	Asp	

945 950 955 960
 5 Ala Gln Leu Phe Leu Arg Phe Gly Gln Val Ile Ser Gln Asn Ser Glu
 965 970 975

 <210> 3
 10 <211> 3000
 <212> DNA
 <213> rice ERECTA homolog
 15

 <220>
 20 <221> CDS
 <222> (1)..(2997)
 <223>
 25

 <400> 3
 30 atg gcg gcg gcg agg gcg ccg tgg ctg tgg tgg tgg gtg gtg gtg gtt
 48
 Met Ala Ala Ala Arg Ala Pro Trp Leu Trp Trp Trp Val Val Val Val
 1 5 10 15
 35 gtt ggt gtg gcg gtg gcg gag gcg gcc tcc gga gga gga gga ggg gga
 96
 Val Gly Val Ala Val Ala Glu Ala Ala Ser Gly Gly Gly Gly Gly Gly
 20 25 30
 40 gat ggg gag ggg aag gcg ctg atg ggc gtg aag gcc ggt ttc ggg aac
 144
 Asp Gly Glu Gly Lys Ala Leu Met Gly Val Lys Ala Gly Phe Gly Asn
 35 40 45
 45 gcg gcc aac gcg ctc gtc gac tgg gac ggc ggc gcc gac cac tgc gcg
 192
 Ala Ala Asn Ala Leu Val Asp Trp Asp Gly Gly Ala Asp His Cys Ala
 50 55 60
 50 tgg cgc ggc gtc acc tgc gac aac gcc tcc ttc gcc gtc ctc gcc ctg
 240
 Trp Arg Gly Val Thr Cys Asp Asn Ala Ser Phe Ala Val Leu Ala Leu
 65 70 75 80
 55 aac ttg tca aat cta aac cta gga ggt gag atc tcg ccg gcc atc gga
 288
 Asn Leu Ser Asn Leu Asn Leu Gly Gly Glu Ile Ser Pro Ala Ile Gly
 85 90 95

gag ctc aag aat cta cag ttc gtt gat ctc aag ggg aac aag ctc act
 336
 5 Glu Leu Lys Asn Leu Gln Phe Val Asp Leu Lys Gly Asn Lys Leu Thr
 100 105 110

ggc caa atc cca gat gag att ggg gac tgc atc tcc tta aaa tat ttg
 384
 10 Gly Gln Ile Pro Asp Glu Ile Gly Asp Cys Ile Ser Leu Lys Tyr Leu
 115 120 125

gat ttg tct ggc aac ttg ctg tat gga gac atc ccc ttc tcc atc tcc
 432
 15 Asp Leu Ser Gly Asn Leu Leu Tyr Gly Asp Ile Pro Phe Ser Ile Ser
 130 135 140

aag ctc aag cag ctt gag gag ctg att ttg aag aac aac cag ctc acg
 480
 20 Lys Leu Lys Gln Leu Glu Glu Leu Ile Leu Lys Asn Asn Gln Leu Thr
 145 150 155 160

gga ccc atc cct tcc aca ttg tcc caa att cca aat ctc aag aca ttg
 528
 25 Gly Pro Ile Pro Ser Thr Leu Ser Gln Ile Pro Asn Leu Lys Thr Leu
 165 170 175

gac ctg gca cag aac cag ctt aca ggc gat atc cca agg ctc ata tac
 576
 30 Asp Leu Ala Gln Asn Gln Leu Thr Gly Asp Ile Pro Arg Leu Ile Tyr
 180 185 190

tgg aat gaa gtt ctg caa tac cta ggt ttg agg ggt aac tca ctg act
 624
 35 Trp Asn Glu Val Leu Gln Tyr Leu Gly Leu Arg Gly Asn Ser Leu Thr
 195 200 205

gga act ttg tca cct gac atg tgc caa ctg act ggc ctg tgg tac ttt
 672
 40 Gly Thr Leu Ser Pro Asp Met Cys Gln Leu Thr Gly Leu Trp Tyr Phe
 210 215 220

gat gta agg gga aac aat ctc aca ggg acc att cca gag agc ata ggg
 720
 45 Asp Val Arg Gly Asn Asn Leu Thr Gly Thr Ile Pro Glu Ser Ile Gly
 225 230 235 240

aac tgc acc agc ttt gag att ctg gac att tcg tat aac caa atc tct
 768
 50 Asn Cys Thr Ser Phe Glu Ile Leu Asp Ile Ser Tyr Asn Gln Ile Ser
 245 250 255

gga gaa ata cct tac aac ata ggc ttt ctt caa gta gcc aca ctg tca
 816
 55 Gly Glu Ile Pro Tyr Asn Ile Gly Phe Leu Gln Val Ala Thr Leu Ser
 260 265 270

ctt caa gga aat aga ctg act ggg aaa att cca gat gtg att ggc ctg
 864
 Leu Gln Gly Asn Arg Leu Thr Gly Lys Ile Pro Asp Val Ile Gly Leu
 275 280 285
 5
 atg caa gct ctt gct gtt cta gac ctg agt gag aac gag ctg gta ggg
 912
 Met Gln Ala Leu Ala Val Leu Asp Leu Ser Glu Asn Glu Leu Val Gly
 290 295 300
 10
 ccc att cct tct ata ctg ggc aat cta tcc tat act gga aaa cta tat
 960
 Pro Ile Pro Ser Ile Leu Gly Asn Leu Ser Tyr Thr Gly Lys Leu Tyr
 305 310 315 320
 15
 tta cat ggg aac aaa ctt act gga gtc ata ccg ccg gag ctt ggg aac
 1008
 Leu His Gly Asn Lys Leu Thr Gly Val Ile Pro Pro Glu Leu Gly Asn
 325 330 335
 20
 atg agt aaa ctt agc tac cta caa ctg aat gat aat gaa ttg gtg ggc
 1056
 Met Ser Lys Leu Ser Tyr Leu Gln Leu Asn Asp Asn Glu Leu Val Gly
 340 345 350
 25
 aca att cca gca gag ctt ggc aaa ctt gaa gag ctt ttt gaa cta aat
 1104
 Thr Ile Pro Ala Glu Leu Gly Lys Leu Glu Glu Leu Phe Glu Leu Asn
 355 360 365
 30
 ctt gcc aac aac aat ctt caa ggt cct att cct gca aac atc agt tct
 1152
 Leu Ala Asn Asn Asn Leu Gln Gly Pro Ile Pro Ala Asn Ile Ser Ser
 370 375 380
 35
 tgc act gct cta aac aaa ttc aat gtt tat ggc aat aag cta aat ggt
 1200
 Cys Thr Ala Leu Asn Lys Phe Asn Val Tyr Gly Asn Lys Leu Asn Gly
 385 390 395 400
 40
 tct att cct gct ggt ttc cag aag ttg gag agt ctg act tac ttg aac
 1248
 Ser Ile Pro Ala Gly Phe Gln Lys Leu Glu Ser Leu Thr Tyr Leu Asn
 405 410 415
 45
 cta tct tca aac aat ttc aaa ggc aat att cct tct gag ctt ggt cac
 1296
 Leu Ser Ser Asn Asn Phe Lys Gly Asn Ile Pro Ser Glu Leu Gly His
 420 425 430
 50
 atc atc aac ttg gac aca ttg gat ctt tcc tac aat gaa ttc tct gga
 1344
 Ile Ile Asn Leu Asp Thr Leu Asp Leu Ser Tyr Asn Glu Phe Ser Gly
 435 440 445
 55
 cca gtt cct gct acc att ggt gat cta gag cac ctt ctt gaa ctg aat
 1392

	Pro	Val	Pro	Ala	Thr	Ile	Gly	Asp	Leu	Glu	His	Leu	Glu	Leu	Glu	Leu	Asn
	450						455					460					
5	ttg	agt	aag	aac	cat	ctt	gat	ggg	cca	gtt	cct	gct	gag	ttt	gga	aac	
	1440																
	Leu	Ser	Lys	Asn	His	Leu	Asp	Gly	Pro	Val	Pro	Ala	Glu	Phe	Gly	Asn	
	465					470					475					480	
10	ttg	aga	agc	gtc	caa	gta	att	gat	atg	tcc	aac	aac	aac	tta	tct	ggt	
	1488																
	Leu	Arg	Ser	Val	Gln	Val	Ile	Asp	Met	Ser	Asn	Asn	Asn	Leu	Ser	Gly	
					485					490					495		
15	agt	ctg	ccc	gag	gaa	ctt	gga	caa	ctt	caa	aac	ctt	gat	agc	ctg	att	
	1536																
	Ser	Leu	Pro	Glu	Glu	Leu	Gly	Gln	Leu	Gln	Asn	Leu	Asp	Ser	Leu	Ile	
				500					505					510			
20	ctt	aac	aac	aac	aat	ttg	gtt	ggg	gag	atc	cct	gct	caa	ttg	gcc	aac	
	1584																
	Leu	Asn	Asn	Asn	Asn	Leu	Val	Gly	Glu	Ile	Pro	Ala	Gln	Leu	Ala	Asn	
				515				520					525				
25	tgc	ttc	agc	tta	aat	aac	ctt	gca	ttt	cag	gaa	ttt	gtc	ata	caa	caa	
	1632																
	Cys	Phe	Ser	Leu	Asn	Asn	Leu	Ala	Phe	Gln	Glu	Phe	Val	Ile	Gln	Gln	
		530					535					540					
30	ttt	atc	tgg	aca	tgt	ccc	gat	ggc	aaa	gaa	ctt	ctc	gaa	att	ccc	aat	
	1680																
	Phe	Ile	Trp	Thr	Cys	Pro	Asp	Gly	Lys	Glu	Leu	Leu	Glu	Ile	Pro	Asn	
	545					550					555					560	
35	gga	aag	cat	ctt	cta	att	tct	gat	tgc	aac	cag	tac	ata	aat	cat	aaa	
	1728																
	Gly	Lys	His	Leu	Leu	Ile	Ser	Asp	Cys	Asn	Gln	Tyr	Ile	Asn	His	Lys	
					565					570					575		
40	tgc	agc	ttc	ttg	ggt	aat	cca	tta	ctg	cat	gtt	tac	tgc	caa	gat	tcc	
	1776																
	Cys	Ser	Phe	Leu	Gly	Asn	Pro	Leu	Leu	His	Val	Tyr	Cys	Gln	Asp	Ser	
				580					585					590			
45	agc	tgt	gga	cac	tct	cat	gga	caa	aga	gtt	aat	att	tca	aag	aca	gca	
	1824																
	Ser	Cys	Gly	His	Ser	His	Gly	Gln	Arg	Val	Asn	Ile	Ser	Lys	Thr	Ala	
			595					600					605				
50	att	gct	tgc	att	atc	tta	ggc	ttt	atc	ata	ttg	ctc	tgc	gtt	ctg	ctg	
	1872																
	Ile	Ala	Cys	Ile	Ile	Leu	Gly	Phe	Ile	Ile	Leu	Le					

gat aag cca gtg caa gga cct cca aag cta gtt gtt ctc cag atg gac
 1968
 5 Asp Lys Pro Val Gln Gly Pro Pro Lys Leu Val Val Leu Gln Met Asp
 645 650 655

atg gct atc cat act tac gag gac atc atg agg ctg aca gag aat ttg
 2016
 10 Met Ala Ile His Thr Tyr Glu Asp Ile Met Arg Leu Thr Glu Asn Leu
 660 665 670

agc gag aaa tac atc att ggc tat ggc gcc tca agc act gtc tac aaa
 2064
 15 Ser Glu Lys Tyr Ile Ile Gly Tyr Gly Ala Ser Ser Thr Val Tyr Lys
 675 680 685

tgt gaa ctc aag agc ggc aag gcc att gct gtc aag cgg ctt tac agt
 2112
 20 Cys Glu Leu Lys Ser Gly Lys Ala Ile Ala Val Lys Arg Leu Tyr Ser
 690 695 700

cag tat aac cat agc ctc cga gag ttt gaa aca gaa cta gag aca att
 2160
 25 Gln Tyr Asn His Ser Leu Arg Glu Phe Glu Thr Glu Leu Glu Thr Ile
 705 710 715 720

ggc agc ata cgg cac agg aat ctt gtt agc ctc cat ggc ttc tcg cta
 2208
 30 Gly Ser Ile Arg His Arg Asn Leu Val Ser Leu His Gly Phe Ser Leu
 725 730 735

tct cca cat gga aac ttg ctc ttc tat gat tac atg gaa aat ggt tcc
 2256
 35 Ser Pro His Gly Asn Leu Leu Phe Tyr Asp Tyr Met Glu Asn Gly Ser
 740 745 750

ttg tgg gat ctt ctc cac ggt cca tca aag aaa gtg aag ctc aac tgg
 2304
 40 Leu Trp Asp Leu Leu His Gly Pro Ser Lys Lys Val Lys Leu Asn Trp
 755 760 765

gac aca aga ctg agg atc gcg gtc gga gct gca caa ggg ctg gcc tat
 2352
 45 Asp Thr Arg Leu Arg Ile Ala Val Gly Ala Ala Gln Gly Leu Ala Tyr
 770 775 780

ctc cac cat gac tgc aac cct cgc ata atc cac aga gat gtc aag tcc
 2400
 50 Leu His His Asp Cys Asn Pro Arg Ile Ile His Arg Asp Val Lys Ser
 785 790 795 800

tcc aac atc ctg ctc gac gag aac ttc gaa gcg cac ctc tca gat ttc
 2448
 55 Ser Asn Ile Leu Leu Asp Glu Asn Phe Glu Ala His Leu Ser Asp Phe
 805 810 815

ggc ata gcc aaa tgt gtc ccc tct gcc aag tcc cat gcc tcc act tat
 2496
 Gly Ile Ala Lys Cys Val Pro Ser Ala Lys Ser His Ala Ser Thr Tyr
 820 825 830

5

gtg cta gga acc atc ggc tac att gat ccg gag tat gcc agg act tcc
 2544
 Val Leu Gly Thr Ile Gly Tyr Ile Asp Pro Glu Tyr Ala Arg Thr Ser
 835 840 845

10

agg ctc aat gag aaa tct gat gtg tac agc ttc ggc atc gtc ctt ctg
 2592
 Arg Leu Asn Glu Lys Ser Asp Val Tyr Ser Phe Gly Ile Val Leu Leu
 850 855 860

15

gaa ttg ctc aca ggg aag aag gcc gtc gac aac gaa tcg aac ttg cat
 2640
 Glu Leu Leu Thr Gly Lys Lys Ala Val Asp Asn Glu Ser Asn Leu His
 865 870 875 880

20

caa ttg ata ctc tcc aaa gct gat gac aac aca gtc atg gag gca gtg
 2688
 Gln Leu Ile Leu Ser Lys Ala Asp Asp Asn Thr Val Met Glu Ala Val
 885 890 895

25

gac tcg gag gtg tca gtg acg tgc acg gac atg gga ctg gtc agg aag
 2736
 Asp Ser Glu Val Ser Val Thr Cys Thr Asp Met Gly Leu Val Arg Lys
 900 905 910

30

gcc ttc cag ctc gcc ctt ctg tgc acc aag agg cac cct tca gac cgg
 2784
 Ala Phe Gln Leu Ala Leu Leu Cys Thr Lys Arg His Pro Ser Asp Arg
 915 920 925

35

ccg acc atg cac gag gtt gca agg gtg ctg ctc tcc ctg ctg ccg gcc
 2832
 Pro Thr Met His Glu Val Ala Arg Val Leu Leu Ser Leu Leu Pro Ala
 930 935 940

40

tcc gcc atg aca acg ccc aag acg gtg gac tac tcc cgg ttg ctg gcg
 2880
 Ser Ala Met Thr Thr Pro Lys Thr Val Asp Tyr Ser Arg Leu Leu Ala
 945 950 955 960

45

tcg acg acg acg gcg gcc gac atg cga ggg cac gac gtg acc gac atc
 2928
 Ser Thr Thr Thr Ala Ala Asp Met Arg Gly His Asp Val Thr Asp Ile
 965 970 975

50

ggc gac aac agc tcc tcc gac gag cag tgg ttc gtc agg ttc ggc gag
 2976
 Gly Asp Asn Ser Ser Ser Asp Glu Gln Trp Phe Val Arg Phe Gly Glu
 980 985 990

55

gtc ata tcc aag cac aca atg tga
 3000

Val Ile Ser Lys His Thr Met
995

5 <210> 4

<211> 999

<212> PRT

10 <213> rice ERECTA homolog

15 <400> 4

Met Ala Ala Ala Arg Ala Pro Trp Leu Trp Trp Trp Val Val Val Val
1 5 10 15

20 Val Gly Val Ala Val Ala Glu Ala Ala Ser Gly Gly Gly Gly Gly Gly
20 25 30

25 Asp Gly Glu Gly Lys Ala Leu Met Gly Val Lys Ala Gly Phe Gly Asn
35 40 45

30 Ala Ala Asn Ala Leu Val Asp Trp Asp Gly Gly Ala Asp His Cys Ala
50 55 60

35 Trp Arg Gly Val Thr Cys Asp Asn Ala Ser Phe Ala Val Leu Ala Leu
65 70 75 80

Asn Leu Ser Asn Leu Asn Leu Gly Gly Glu Ile Ser Pro Ala Ile Gly
85 90 95

40 Glu Leu Lys Asn Leu Gln Phe Val Asp Leu Lys Gly Asn Lys Leu Thr
100 105 110

45 Gly Gln Ile Pro Asp Glu Ile Gly Asp Cys Ile Ser Leu Lys Tyr Leu
115 120 125

50 Asp Leu Ser Gly Asn Leu Leu Tyr Gly Asp Ile Pro Phe Ser Ile Ser
130 135 140

55 Lys Leu Lys Gln Leu Glu Glu Leu Ile Leu Lys Asn Asn Gln Leu Thr
145 150 155 160

Gly Pro Ile Pro Ser Thr Leu Ser Gln Ile Pro Asn Leu Lys Thr Leu

	165	170	175
5	Asp Leu Ala Gln Asn Gln Leu Thr Gly Asp Ile Pro Arg Leu Ile Tyr 180 185 190		
10	Trp Asn Glu Val Leu Gln Tyr Leu Gly Leu Arg Gly Asn Ser Leu Thr 195 200 205		
15	Gly Thr Leu Ser Pro Asp Met Cys Gln Leu Thr Gly Leu Trp Tyr Phe 210 215 220		
20	Asp Val Arg Gly Asn Asn Leu Thr Gly Thr Ile Pro Glu Ser Ile Gly 225 230 235 240		
25	Asn Cys Thr Ser Phe Glu Ile Leu Asp Ile Ser Tyr Asn Gln Ile Ser 245 250 255		
30	Gly Glu Ile Pro Tyr Asn Ile Gly Phe Leu Gln Val Ala Thr Leu Ser 260 265 270		
35	Leu Gln Gly Asn Arg Leu Thr Gly Lys Ile Pro Asp Val Ile Gly Leu 275 280 285		
40	Met Gln Ala Leu Ala Val Leu Asp Leu Ser Glu Asn Glu Leu Val Gly 290 295 300		
45	Pro Ile Pro Ser Ile Leu Gly Asn Leu Ser Tyr Thr Gly Lys Leu Tyr 305 310 315 320		
50	Leu His Gly Asn Lys Leu Thr Gly Val Ile Pro Pro Glu Leu Gly Asn 325 330 335		
55	Met Ser Lys Leu Ser Tyr Leu Gln Leu Asn Asp Asn Glu Leu Val Gly 340 345 350		
60	Thr Ile Pro Ala Glu Leu Gly Lys Leu Glu Glu Leu Phe Glu Leu Asn 355 360 365		
65	Leu Ala Asn Asn Asn Leu Gln Gly Pro Ile Pro Ala Asn Ile Ser Ser 370 375 380		
70	Cys Thr Ala Leu Asn Lys Phe Asn Val Tyr Gly Asn Lys Leu Asn Gly 385 390 395 400		

Ser Ile Pro Ala Gly Phe Gln Lys Leu Glu Ser Leu Thr Tyr Leu Asn
 405 410 415
 5

Leu Ser Ser Asn Asn Phe Lys Gly Asn Ile Pro Ser Glu Leu Gly His
 420 425 430

10

Ile Ile Asn Leu Asp Thr Leu Asp Leu Ser Tyr Asn Glu Phe Ser Gly
 435 440 445

15

Pro Val Pro Ala Thr Ile Gly Asp Leu Glu His Leu Leu Glu Leu Asn
 450 455 460

20

Leu Ser Lys Asn His Leu Asp Gly Pro Val Pro Ala Glu Phe Gly Asn
 465 470 475 480

25

Leu Arg Ser Val Gln Val Ile Asp Met Ser Asn Asn Asn Leu Ser Gly
 485 490 495

30

Ser Leu Pro Glu Glu Leu Gly Gln Leu Gln Asn Leu Asp Ser Leu Ile
 500 505 510

35

Leu Asn Asn Asn Asn Leu Val Gly Glu Ile Pro Ala Gln Leu Ala Asn
 515 520 525

40

Cys Phe Ser Leu Asn Asn Leu Ala Phe Gln Glu Phe Val Ile Gln Gln
 530 535 540

45

Phe Ile Trp Thr Cys Pro Asp Gly Lys Glu Leu Leu Glu Ile Pro Asn
 545 550 555 560

Gly Lys His Leu Leu Ile Ser Asp Cys Asn Gln Tyr Ile Asn His Lys
 565 570 575

50

Cys Ser Phe Leu Gly Asn Pro Leu Leu His Val Tyr Cys Gln Asp Ser
 580 585 590

55

Ser Cys Gly His Ser His Gly Gln Arg Val Asn Ile Ser Lys Thr Ala
 595 600 605

Ile Ala Cys Ile Ile Leu Gly Phe Ile Ile Leu Leu Cys Val Leu Leu
 610 615 620

Leu Ala Ile Tyr Lys Thr Asn Gln Pro Gln Pro Leu Val Lys Gly Ser
 625 630 635 640

5 Asp Lys Pro Val Gln Gly Pro Pro Lys Leu Val Val Leu Gln Met Asp
 645 650 655

10 Met Ala Ile His Thr Tyr Glu Asp Ile Met Arg Leu Thr Glu Asn Leu
 660 665 670

15 Ser Glu Lys Tyr Ile Ile Gly Tyr Gly Ala Ser Ser Thr Val Tyr Lys
 675 680 685

20 Cys Glu Leu Lys Ser Gly Lys Ala Ile Ala Val Lys Arg Leu Tyr Ser
 690 695 700

Gln Tyr Asn His Ser Leu Arg Glu Phe Glu Thr Glu Leu Glu Thr Ile
 705 710 715 720

25 Gly Ser Ile Arg His Arg Asn Leu Val Ser Leu His Gly Phe Ser Leu
 725 730 735

30 Ser Pro His Gly Asn Leu Leu Phe Tyr Asp Tyr Met Glu Asn Gly Ser
 740 745 750

35 Leu Trp Asp Leu Leu His Gly Pro Ser Lys Lys Val Lys Leu Asn Trp
 755 760 765

40 Asp Thr Arg Leu Arg Ile Ala Val Gly Ala Ala Gln Gly Leu Ala Tyr
 770 775 780

Leu His His Asp Cys Asn Pro Arg Ile Ile His Arg Asp Val Lys Ser
 785 790 795 800

45 Ser Asn Ile Leu Leu Asp Glu Asn Phe Glu Ala His Leu Ser Asp Phe
 805 810 815

50 Gly Ile Ala Lys Cys Val Pro Ser Ala Lys Ser His Ala Ser Thr Tyr
 820 825 830

55 Val Leu Gly Thr Ile Gly Tyr Ile Asp Pro Glu Tyr Ala Arg Thr Ser
 835 840 845

Arg Leu Asn Glu Lys Ser Asp Val Tyr Ser Phe Gly Ile Val Leu Leu
 850 855 860

5 Glu Leu Leu Thr Gly Lys Lys Ala Val Asp Asn Glu Ser Asn Leu His
 865 870 875 880

10 Gln Leu Ile Leu Ser Lys Ala Asp Asp Asn Thr Val Met Glu Ala Val
 885 890 895

15 Asp Ser Glu Val Ser Val Thr Cys Thr Asp Met Gly Leu Val Arg Lys
 900 905 910

20 Ala Phe Gln Leu Ala Leu Leu Cys Thr Lys Arg His Pro Ser Asp Arg
 915 920 925

25 Ser Ala Met Thr Thr Pro Lys Thr Val Asp Tyr Ser Arg Leu Leu Ala
 945 950 955 960

30 Ser Thr Thr Thr Ala Ala Asp Met Arg Gly His Asp Val Thr Asp Ile
 965 970 975

35 Gly Asp Asn Ser Ser Ser Asp Glu Gln Trp Phe Val Arg Phe Gly Glu
 980 985 990

Val Ile Ser Lys His Thr Met
 995

40 <210> 5
 <211> 2766

45 <212> DNA
 <213> sorghum ERECTA homolog

50 <220>
 <221> CDS

55 <222> (1)..(2763)
 <223>

```

<400> 5
5  atg acg acg acg gcc gcc cgt gct ctc gtc gcc ctc ctc ctc gtc gcc
   48
   Met Thr Thr Thr Ala Ala Arg Ala Leu Val Ala Leu Leu Leu Val Ala
   1           5           10           15

10  gtc gcc gtc gcc gac gat ggg gcg acg ctg gtg gag atc aag aag tcc
   96
   Val Ala Val Ala Asp Asp Gly Ala Thr Leu Val Glu Ile Lys Lys Ser
           20           25           30

15  ttc cgc aac gtc ggc aac gta ctg tac gat tgg gcc ggc gac gac tac
   144
   Phe Arg Asn Val Gly Asn Val Leu Tyr Asp Trp Ala Gly Asp Asp Tyr
           35           40           45

20  tgc tcc tgg cgc ggc gtc ctg tgc gac aac gtc aca ttc gcc gtc gct
   192
   Cys Ser Trp Arg Gly Val Leu Cys Asp Asn Val Thr Phe Ala Val Ala
   50           55           60

25  gcg ctc aac ctc tct ggc ctc aac ctt gag ggc gag atc tct cca gcc
   240
   Ala Leu Asn Leu Ser Gly Leu Asn Leu Glu Gly Glu Ile Ser Pro Ala
   65           70           75           80

30  gtc ggc agc ctc aag agc ctc gtc tcc atc gat ctg aag tca aat ggg
   288
   Val Gly Ser Leu Lys Ser Leu Val Ser Ile Asp Leu Lys Ser Asn Gly
           85           90           95

35  cta tcc ggg cag atc cct gat gag att ggt gat tgt tca tca ctt agg
   336
   Leu Ser Gly Gln Ile Pro Asp Glu Ile Gly Asp Cys Ser Ser Leu Arg
           100           105           110

40  acg ctg gac ttt tct ttc aac aac ttg gat ggc gac ata cca ttt tct
   384
   Thr Leu Asp Phe Ser Phe Asn Asn Leu Asp Gly Asp Ile Pro Phe Ser
           115           120           125

45  ata tca aag ctg aag cac ctg gag aac ttg ata ttg aag aac aac cag
   432
   Ile Ser Lys Leu Lys His Leu Glu Asn Leu Ile Leu Lys Asn Asn Gln
   130           135           140

50  ctg att ggt gcg atc cca tca aca ttg tca cag ctc cca aat ttg aag
   480
   Leu Ile Gly Ala Ile Pro Ser Thr Leu Ser Gln Leu Pro Asn Leu Lys
   145           150           155           160

55  att ttg gat ttg gca caa aac aaa ctg act ggg gag ata cca agg ctt
   528
   Ile Leu Asp Leu Ala Gln Asn Lys Leu Thr Gly Glu Ile Pro Arg Leu

```

	165		170		175
	atc tac tgg aat gag gtt ctt caa tat ctt gat gtg aag aac aat agc				
	576				
5	Ile Tyr Trp Asn Glu Val Leu Gln Tyr Leu Asp Val Lys Asn Asn Ser				
		180		185	190
	ttg acc ggg gtg ata cca gac acc att ggg aac tgt aca agt ttt caa				
	624				
10	Leu Thr Gly Val Ile Pro Asp Thr Ile Gly Asn Cys Thr Ser Phe Gln				
		195		200	205
	gtc ttg gat ttg tct tac aac cgc ttt act gga cca atc cca ttc aac				
	672				
15	Val Leu Asp Leu Ser Tyr Asn Arg Phe Thr Gly Pro Ile Pro Phe Asn				
		210		215	220
	att ggt ttc cta caa gtg gct aca cta tcc ttg caa ggg aac aag ttc				
	720				
20	Ile Gly Phe Leu Gln Val Ala Thr Leu Ser Leu Gln Gly Asn Lys Phe				
		225		230	235
	acc ggt cca att cct tca gta att ggt ctt atg cag gct ctc gct gtt				
	768				
25	Thr Gly Pro Ile Pro Ser Val Ile Gly Leu Met Gln Ala Leu Ala Val				
		245		250	255
	cta gat ctg agt tac aac caa tta tct ggt cct ata cca tca ata cta				
	816				
30	Leu Asp Leu Ser Tyr Asn Gln Leu Ser Gly Pro Ile Pro Ser Ile Leu				
		260		265	270
	ggc aac ttg aca tac act gag aag ctg tac atc caa ggc aat aag tta				
	864				
35	Gly Asn Leu Thr Tyr Thr Glu Lys Leu Tyr Ile Gln Gly Asn Lys Leu				
		275		280	285
	act ggg tgg ata cca cca gag tta gga aat atg tca aca ctt cat tac				
	912				
40	Thr Gly Ser Ile Pro Pro Glu Leu Gly Asn Met Ser Thr Leu His Tyr				
		290		295	300
	cta gaa ctg aac gat aat caa ctt act ggg tca att cca cca gag ctt				
	960				
45	Leu Glu Leu Asn Asp Asn Gln Leu Thr Gly Ser Ile Pro Pro Glu Leu				
		305		310	315
	gga agg cta aca ggc ttg ttt gac ctg aac ctt gcg aat aac cac ctg				
	1008				
50	Gly Arg Leu Thr Gly Leu Phe Asp Leu Asn Leu Ala Asn Asn His Leu				
		325		330	335
	gaa gga cca att cct gac aac cta agt tca tgt gtg aat ctc aat agc				
	1056				
55	Glu Gly Pro Ile Pro Asp Asn Leu Ser Ser Cys Val Asn Leu Asn Ser				
		340		345	350

ttc aat gct tat ggc aac aag tta aat ggg acc att cct cgt tcg ttg
 1104
 Phe Asn Ala Tyr Gly Asn Lys Leu Asn Gly Thr Ile Pro Arg Ser Leu
 355 360 365
 5
 cgg aaa ctt gaa agc atg acc tat tta aat ctg tca tca aac ttc ata
 1152
 Arg Lys Leu Glu Ser Met Thr Tyr Leu Asn Leu Ser Ser Asn Phe Ile
 370 375 380
 10
 agt ggc tct att cct att gag tta tca agg atc aac aat ttg gac acg
 1200
 Ser Gly Ser Ile Pro Ile Glu Leu Ser Arg Ile Asn Asn Leu Asp Thr
 385 390 395 400
 15
 ctg gat tta tcc tgt aac atg atg act ggt cca att cca tca tca att
 1248
 Leu Asp Leu Ser Cys Asn Met Met Thr Gly Pro Ile Pro Ser Ser Ile
 405 410 415
 20
 ggc agc cta gag cat cta ttg aga ctt aac ttg agc aag aat ggt cta
 1296
 Gly Ser Leu Glu His Leu Leu Arg Leu Asn Leu Ser Lys Asn Gly Leu
 420 425 430
 25
 gtt gga ttc atc ccc gcg gag ttt ggt aat ttg agg agt gtc atg gag
 1344
 Val Gly Phe Ile Pro Ala Glu Phe Gly Asn Leu Arg Ser Val Met Glu
 435 440 445
 30
 att gat tta tcc tat aat cac ctt ggt ggc ctg att cct caa gaa ctt
 1392
 Ile Asp Leu Ser Tyr Asn His Leu Gly Gly Leu Ile Pro Gln Glu Leu
 450 455 460
 35
 gaa atg ctg caa aac ctg atg ttg cta aat gtg tcg tac aat aat ttg
 1440
 Glu Met Leu Gln Asn Leu Met Leu Leu Asn Val Ser Tyr Asn Asn Leu
 465 470 475 480
 40
 gct ggt gtt gtc cct gct gac aac aac ttc aca cgg ttt tca cct gac
 1488
 Ala Gly Val Val Pro Ala Asp Asn Asn Phe Thr Arg Phe Ser Pro Asp
 485 490 495
 45
 agc ttt tta ggt aat cct gga ctc tgt gga tac tgg ctt ggt tcg tcg
 1536
 Ser Phe Leu Gly Asn Pro Gly Leu Cys Gly Tyr Trp Leu Gly Ser Ser
 500 505 510
 50
 tgt cgt tcc act ggc cac cac gag aaa ccg cct atc tca aag gct gcc
 1584
 Cys Arg Ser Thr Gly His His Glu Lys Pro Pro Ile Ser Lys Ala Ala
 515 520 525
 55
 ata att ggt gtt gct gtg ggt gga ctt gtt atc ctc ttg atg atc tta
 1632

	Ile	Ile	Gly	Val	Ala	Val	Gly	Gly	Leu	Val	Ile	Leu	Leu	Met	Ile	Leu
	530						535					540				
5	gta	gct	gtt	tgc	agg	cca	cat	cgt	cca	cct	gct	ttt	aaa	gat	gtc	act
	1680															
	Val	Ala	Val	Cys	Arg	Pro	His	Arg	Pro	Pro	Ala	Phe	Lys	Asp	Val	Thr
	545					550					555					560
10	gta	agc	aag	cca	gtg	aga	aat	gct	ccc	ccc	aag	ctg	gtg	atc	ctt	cat
	1728															
	Val	Ser	Lys	Pro	Val	Arg	Asn	Ala	Pro	Pro	Lys	Leu	Val	Ile	Leu	His
					565					570					575	
15	atg	aac	atg	gcc	ctt	cat	gta	tac	gat	gac	ata	atg	agg	atg	act	gag
	1776															
	Met	Asn	Met	Ala	Leu	His	Val	Tyr	Asp	Asp	Ile	Met	Arg	Met	Thr	Glu
				580					585					590		
20	aac	ttg	agt	gag	aaa	tac	atc	att	gga	tac	ggg	gcg	tca	agt	aca	gtt
	1824															
	Asn	Leu	Ser	Glu	Lys	Tyr	Ile	Ile	Gly	Tyr	Gly	Ala	Ser	Ser	Thr	Val
			595					600					605			
25	tat	aaa	tgt	gtc	cta	aag	aat	tgc	aaa	ccg	gtg	gca	ata	aaa	aag	ctg
	1872															
	Tyr	Lys	Cys	Val	Leu	Lys	Asn	Cys	Lys	Pro	Val	Ala	Ile	Lys	Lys	Leu
		610					615					620				
30	tat	gcc	cac	tac	cca	cag	agc	ctt	aag	gaa	ttt	gaa	act	gag	ctt	gag
	1920															
	Tyr	Ala	His	Tyr	Pro	Gln	Ser	Leu	Lys	Glu	Phe	Glu	Thr	Glu	Leu	Glu
		625				630					635					640
35	act	gtt	ggt	agc	atc	aag	cac	cgg	aat	cta	gtc	agc	ctt	caa	ggg	tac
	1968															
	Thr	Val	Gly	Ser	Ile	Lys	His	Arg	Asn	Leu	Val	Ser	Leu	Gln	Gly	Tyr
					645					650					655	
40	tca	tta	tca	cct	gtt	ggg	aac	ctc	ctc	ttt	tat	gat	tat	atg	gaa	tgt
	2016															
	Ser	Leu	Ser	Pro	Val	Gly	Asn	Leu	Leu	Phe	Tyr	Asp	Tyr	Met	Glu	Cys
				660					665					670		
45	ggc	agc	tta	tgg	gat	gtt	tta	cat	gaa	ggt	tca	tcc	aag	aag	aaa	aaa
	2064															
	Gly	Ser	Leu	Trp	Asp	Val	Leu	His	Glu	Gly	Ser	Ser	Lys	Lys	Lys	Lys
			675					680					685			
50	ctt	gac	tgg	gag	act	cgc	cta	cgg	att	gct	ctt	ggt	gca	gct	caa	ggc
	2112															
	Leu	Asp	Trp	Glu	Thr	Arg	Leu	Arg	Ile	Ala	Leu	Gly	Ala	Ala	Gln	Gly
		690					695					700				
55	ctt	gct	tac	ctt	cac	cat	gac	tgc	agt	cca	cgg	ata	att	cat	cgg	gat
	2160															
	Leu	Ala	Tyr	Leu	His	His	Asp	Cys	Ser	Pro	Arg	Ile	Ile	His	Arg	Asp
		705				710					715					720

	gta aaa tca aag aat ata ctc ctt gac aaa gat tat gag gcc cat ctt	
	2208	
5	Val Lys Ser Lys Asn Ile Leu Leu Asp Lys Asp Tyr Glu Ala His Leu	725 730 735
	aca gac ttt gga att gct aag agc tta tgt gtc tca aaa act cac aca	
	2256	
10	Thr Asp Phe Gly Ile Ala Lys Ser Leu Cys Val Ser Lys Thr His Thr	740 745 750
	tca acc tat gtc atg gga act att ggc tac att gat cct gag tac gcc	
	2304	
15	Ser Thr Tyr Val Met Gly Thr Ile Gly Tyr Ile Asp Pro Glu Tyr Ala	755 760 765
	cgc act tcc cgt ctc aac gaa aag tct gat gtc tac agg cta tgg cat	
	2352	
20	Arg Thr Ser Arg Leu Asn Glu Lys Ser Asp Val Tyr Arg Leu Trp His	770 775 780
	tgt tct gct gga gct gct gac tgg caa gaa gcc agt gga caa cga atc	
	2400	
25	Cys Ser Ala Gly Ala Ala Asp Trp Gln Glu Ala Ser Gly Gln Arg Ile	785 790 795 800
	cta tcg aag acg gca agc aac gag gtc atg gat acc gtg gac cct gac	
	2448	
30	Leu Ser Lys Thr Ala Ser Asn Glu Val Met Asp Thr Val Asp Pro Asp	805 810 815
	atc ggg gac acc tgc aag gac ctc ggc gag gtg aag aag ctg ttc cag	
	2496	
35	Ile Gly Asp Thr Cys Lys Asp Leu Gly Glu Val Lys Lys Leu Phe Gln	820 825 830
	ctg gcg ctc ctt tgc acc aag cgg caa ccc tcg gac cga ccg acg atg	
	2544	
40	Leu Ala Leu Leu Cys Thr Lys Arg Gln Pro Ser Asp Arg Pro Thr Met	835 840 845
	cac gag gtg gtg cgc gtc ctg gac tgc ctg gtg aac ccg gac ccg ccg	
	2592	
45	His Glu Val Val Arg Val Leu Asp Cys Leu Val Asn Pro Asp Pro Pro	850 855 860
	cca aag ccg tcg gcg cac cag ctg ccg cag ccg tcg cca gcc gtg cca	
	2640	
50	Pro Lys Pro Ser Ala His Gln Leu Pro Gln Pro Ser Pro Ala Val Pro	865 870 875 880
	agc tac atc aac gag tac gtc agc ctg cgg ggc acc ggc gct ctc tcc	
	2688	
55	Ser Tyr Ile Asn Glu Tyr Val Ser Leu Arg Gly Thr Gly Ala Leu Ser	885 890 895

tgc gcc aac tcg acc agc acc tcg gac gcc gag ctg ttc ctc aag ttc
 2736
 Cys Ala Asn Ser Thr Ser Thr Ser Asp Ala Glu Leu Phe Leu Lys Phe
 900 905 910

5

ggc gag gcc atc tcg cag aac atg gag tag
 2766
 Gly Glu Ala Ile Ser Gln Asn Met Glu
 915 920

10

<210> 6

<211> 921

15

<212> PRT

<213> sorghum ERECTA homolog

20

<400> 6

Met Thr Thr Thr Ala Ala Arg Ala Leu Val Ala Leu Leu Leu Val Ala
 25 1 5 10 15

Val Ala Val Ala Asp Asp Gly Ala Thr Leu Val Glu Ile Lys Lys Ser
 20 25 30

30

Phe Arg Asn Val Gly Asn Val Leu Tyr Asp Trp Ala Gly Asp Asp Tyr
 35 40 45

35

Cys Ser Trp Arg Gly Val Leu Cys Asp Asn Val Thr Phe Ala Val Ala
 50 55 60

40

Ala Leu Asn Leu Ser Gly Leu Asn Leu Glu Gly Glu Ile Ser Pro Ala
 65 70 75 80

45

Val Gly Ser Leu Lys Ser Leu Val Ser Ile Asp Leu Lys Ser Asn Gly
 85 90 95

Leu Ser Gly Gln Ile Pro Asp Glu Ile Gly Asp Cys Ser Ser Leu Arg
 100 105 110

50

Thr Leu Asp Phe Ser Phe Asn Asn Leu Asp Gly Asp Ile Pro Phe Ser
 115 120 125

55

Ile Ser Lys Leu Lys His Leu Glu Asn Leu Ile Leu Lys Asn Asn Gln
 130 135 140

Leu Ile Gly Ala Ile Pro Ser Thr Leu Ser Gln Leu Pro Asn Leu Lys
 145 150 155 160
 5

Ile Leu Asp Leu Ala Gln Asn Lys Leu Thr Gly Glu Ile Pro Arg Leu
 165 170 175

10

Ile Tyr Trp Asn Glu Val Leu Gln Tyr Leu Asp Val Lys Asn Asn Ser
 180 185 190

15

Leu Thr Gly Val Ile Pro Asp Thr Ile Gly Asn Cys Thr Ser Phe Gln
 195 200 205

20

Val Leu Asp Leu Ser Tyr Asn Arg Phe Thr Gly Pro Ile Pro Phe Asn
 210 215 220

25

Ile Gly Phe Leu Gln Val Ala Thr Leu Ser Leu Gln Gly Asn Lys Phe
 225 230 235 240

Thr Gly Pro Ile Pro Ser Val Ile Gly Leu Met Gln Ala Leu Ala Val
 245 250 255

30

Leu Asp Leu Ser Tyr Asn Gln Leu Ser Gly Pro Ile Pro Ser Ile Leu
 260 265 270

35

Gly Asn Leu Thr Tyr Thr Glu Lys Leu Tyr Ile Gln Gly Asn Lys Leu
 275 280 285

40

Thr Gly Ser Ile Pro Pro Glu Leu Gly Asn Met Ser Thr Leu His Tyr
 290 295 300

45

Leu Glu Leu Asn Asp Asn Gln Leu Thr Gly Ser Ile Pro Pro Glu Leu
 305 310 315 320

Gly Arg Leu Thr Gly Leu Phe Asp Leu Asn Leu Ala Asn Asn His Leu
 325 330 335

50

Glu Gly Pro Ile Pro Asp Asn Leu Ser Ser Cys Val Asn Leu Asn Ser
 340 345 350

55

Phe Asn Ala Tyr Gly Asn Lys Leu Asn Gly Thr Ile Pro Arg Ser Leu
 355 360 365

	Arg	Lys	Leu	Glu	Ser	Met	Thr	Tyr	Leu	Asn	Leu	Ser	Ser	Asn	Phe	Ile
	370						375					380				
5	Ser	Gly	Ser	Ile	Pro	Ile	Glu	Leu	Ser	Arg	Ile	Asn	Asn	Leu	Asp	Thr
	385					390					395					400
10	Leu	Asp	Leu	Ser	Cys	Asn	Met	Met	Thr	Gly	Pro	Ile	Pro	Ser	Ser	Ile
					405					410						415
15	Gly	Ser	Leu	Glu	His	Leu	Leu	Arg	Leu	Asn	Leu	Ser	Lys	Asn	Gly	Leu
				420					425					430		
20	Val	Gly	Phe	Ile	Pro	Ala	Glu	Phe	Gly	Asn	Leu	Arg	Ser	Val	Met	Glu
			435					440					445			
	Ile	Asp	Leu	Ser	Tyr	Asn	His	Leu	Gly	Gly	Leu	Ile	Pro	Gln	Glu	Leu
	450						455					460				
25	Glu	Met	Leu	Gln	Asn	Leu	Met	Leu	Leu	Asn	Val	Ser	Tyr	Asn	Asn	Leu
	465					470					475					480
30	Ala	Gly	Val	Val	Pro	Ala	Asp	Asn	Asn	Phe	Thr	Arg	Phe	Ser	Pro	Asp
					485					490						495
35	Ser	Phe	Leu	Gly	Asn	Pro	Gly	Leu	Cys	Gly	Tyr	Trp	Leu	Gly	Ser	Ser
				500					505					510		
40	Cys	Arg	Ser	Thr	Gly	His	His	Glu	Lys	Pro	Pro	Ile	Ser	Lys	Ala	Ala
			515					520					525			
	Ile	Ile	Gly	Val	Ala	Val	Gly	Gly	Leu	Val	Ile	Leu	Leu	Met	Ile	Leu
	530						535					540				
45	Val	Ala	Val	Cys	Arg	Pro	His	Arg	Pro	Pro	Ala	Phe	Lys	Asp	Val	Thr
	545					550					555					560
50	Val	Ser	Lys	Pro	Val	Arg	Asn	Ala	Pro	Pro	Lys	Leu	Val	Ile	Leu	His
					565					570					575	
55	Met	Asn	Met	Ala	Leu	His	Val	Tyr	Asp	Asp	Ile	Met	Arg	Met	Thr	Glu
				580					585					590		

Asn Leu Ser Glu Lys Tyr Ile Ile Gly Tyr Gly Ala Ser Ser Thr Val
 595 600 605

5 Tyr Lys Cys Val Leu Lys Asn Cys Lys Pro Val Ala Ile Lys Lys Leu
 610 615 620

10 Tyr Ala His Tyr Pro Gln Ser Leu Lys Glu Phe Glu Thr Glu Leu Glu
 625 630 635 640

15 Thr Val Gly Ser Ile Lys His Arg Asn Leu Val Ser Leu Gln Gly Tyr
 645 650 655

Ser Leu Ser Pro Val Gly Asn Leu Leu Phe Tyr Asp Tyr Met Glu Cys
 660 665 670

20 Gly Ser Leu Trp Asp Val Leu His Glu Gly Ser Ser Lys Lys Lys Lys
 675 680 685

25 Leu Asp Trp Glu Thr Arg Leu Arg Ile Ala Leu Gly Ala Ala Gln Gly
 690 695 700

30 Leu Ala Tyr Leu His His Asp Cys Ser Pro Arg Ile Ile His Arg Asp
 705 710 715 720

35 Val Lys Ser Lys Asn Ile Leu Leu Asp Lys Asp Tyr Glu Ala His Leu
 725 730 735

40 Thr Asp Phe Gly Ile Ala Lys Ser Leu Cys Val Ser Lys Thr His Thr
 740 745 750

Ser Thr Tyr Val Met Gly Thr Ile Gly Tyr Ile Asp Pro Glu Tyr Ala
 755 760 765

45 Arg Thr Ser Arg Leu Asn Glu Lys Ser Asp Val Tyr Arg Leu Trp His
 770 775 780

50 Cys Ser Ala Gly Ala Ala Asp Trp Gln Glu Ala Ser Gly Gln Arg Ile
 785 790 795 800

55 Leu Ser Lys Thr Ala Ser Asn Glu Val Met Asp Thr Val Asp Pro Asp
 805 810 815

Ile Gly Asp Thr Cys Lys Asp Leu Gly Glu Val Lys Lys Leu Phe Gln

820 825 830
 5 Leu Ala Leu Leu Cys Thr Lys Arg Gln Pro Ser Asp Arg Pro Thr Met
 835 840 845
 10 His Glu Val Val Arg Val Leu Asp Cys Leu Val Asn Pro Asp Pro Pro
 850 855 860
 15 Pro Lys Pro Ser Ala His Gln Leu Pro Gln Pro Ser Pro Ala Val Pro
 865 870 875 880
 20 Ser Tyr Ile Asn Glu Tyr Val Ser Leu Arg Gly Thr Gly Ala Leu Ser
 885 890 895
 25 Cys Ala Asn Ser Thr Ser Thr Ser Asp Ala Glu Leu Phe Leu Lys Phe
 900 905 910
 30 Gly Glu Ala Ile Ser Gln Asn Met Glu
 915 920
 35 <210> 7
 40 <211> 2799
 <212> DNA
 <213> Arabdopsis thaliana ERECTA homolog
 45 <220>
 <221> CDS
 <222> (1)..(2796)
 <223>
 50 <400> 7
 atg gcg ata aag gct tca ttc agc aac gtg gcg aat atg ctt ctt gat
 48
 Met Ala Ile Lys Ala Ser Phe Ser Asn Val Ala Asn Met Leu Leu Asp
 1 5 10 15
 55 tgg gac gat gtt cat aac cac gac ttt tgt tct tgg aga ggt gtc ttc
 96
 Trp Asp Asp Val His Asn His Asp Phe Cys Ser Trp Arg Gly Val Phe
 20 25 30

tgt gat aac gtt agc ctc aat gtt gtc tct ctt aat ctg tca aac ctg
 144
 5 Cys Asp Asn Val Ser Leu Asn Val Val Ser Leu Asn Leu Ser Asn Leu
 35 40 45

aat ctt ggt gga gag ata tca tct gcc ctt gga gat ttg atg aat ctg
 192
 10 Asn Leu Gly Gly Glu Ile Ser Ser Ala Leu Gly Asp Leu Met Asn Leu
 50 55 60

caa tca ata gac ttg caa gga aat aaa ttg ggt ggt caa att cca gat
 240
 15 Gln Ser Ile Asp Leu Gln Gly Asn Lys Leu Gly Gly Gln Ile Pro Asp
 65 70 75 80

gag att gga aac tgt gtt tct ctt gct tat gtg gat ttc tcc acc aat
 288
 20 Glu Ile Gly Asn Cys Val Ser Leu Ala Tyr Val Asp Phe Ser Thr Asn
 85 90 95

ttg ttg ttt gga gac ata ccg ttt tca atc tct aaa ctc aaa cag ctg
 336
 25 Leu Leu Phe Gly Asp Ile Pro Phe Ser Ile Ser Lys Leu Lys Gln Leu
 100 105 110

gag ttt ctg aac cta aag aat aat cag ctc act ggt cca ata cca gca
 384
 30 Glu Phe Leu Asn Leu Lys Asn Asn Gln Leu Thr Gly Pro Ile Pro Ala
 115 120 125

acc tta act cag att cca aac ctt aag acc ctt gac ctc gca aga aac
 432
 35 Thr Leu Thr Gln Ile Pro Asn Leu Lys Thr Leu Asp Leu Ala Arg Asn
 130 135 140

cag ctt act ggt gag ata cca agg tta ctc tac tgg aat gaa gtt tta
 480
 40 Gln Leu Thr Gly Glu Ile Pro Arg Leu Leu Tyr Trp Asn Glu Val Leu
 145 150 155 160

cag tat ctc ggt tta cgt ggg aat atg tta act ggg aca ttg tct cct
 528
 45 Gln Tyr Leu Gly Leu Arg Gly Asn Met Leu Thr Gly Thr Leu Ser Pro
 165 170 175

gat atg tgt cag ctg acg ggt ctg tgg tac ttt gat gtg aga ggc aac
 576
 50 Asp Met Cys Gln Leu Thr Gly Leu Trp Tyr Phe Asp Val Arg Gly Asn
 180 185 190

aac ctt act gga act atc cca gag agc att ggc aat tgc aca agc ttt
 624
 55 Asn Leu Thr Gly Thr Ile Pro Glu Ser Ile Gly Asn Cys Thr Ser Phe
 195 200 205

gag atc ttg gat gta tct tat aat cag att acc gga gtt ata ccc tac
 672
 Glu Ile Leu Asp Val Ser Tyr Asn Gln Ile Thr Gly Val Ile Pro Tyr
 210 215 220
 5
 aat att ggt ttc ctc caa gta gct act ctg tca ctt caa gga aac aag
 720
 Asn Ile Gly Phe Leu Gln Val Ala Thr Leu Ser Leu Gln Gly Asn Lys
 225 230 235 240
 10
 ttg act ggc aga att ccg gaa gtg att ggt ctg atg cag gct ctt gct
 768
 Leu Thr Gly Arg Ile Pro Glu Val Ile Gly Leu Met Gln Ala Leu Ala
 245 250 255
 15
 gta ttg gat ttg agt gac aat gaa tta act ggg cct att cca cca ata
 816
 Val Leu Asp Leu Ser Asp Asn Glu Leu Thr Gly Pro Ile Pro Pro Ile
 260 265 270
 20
 ctt ggg aat ctg tca ttc act gga aaa ctg tat ctc cat ggc aac aag
 864
 Leu Gly Asn Leu Ser Phe Thr Gly Lys Leu Tyr Leu His Gly Asn Lys
 275 280 285
 25
 ctc act gga caa atc cca ccc gag cta ggc aat atg tca cga ctc agc
 912
 Leu Thr Gly Gln Ile Pro Pro Glu Leu Gly Asn Met Ser Arg Leu Ser
 290 295 300
 30
 tat ttg caa cta aat gat aat gaa cta gtg gga aag atc cca cct gag
 960
 Tyr Leu Gln Leu Asn Asp Asn Glu Leu Val Gly Lys Ile Pro Pro Glu
 305 310 315 320
 35
 ctt ggg aag ctg gaa caa ttg ttc gaa ctg aat ctt gcg aac aac aat
 1008
 Leu Gly Lys Leu Glu Gln Leu Phe Glu Leu Asn Leu Ala Asn Asn Asn
 325 330 335
 40
 ctt gta ggg ctg att cca tct aac att agt tcc tgt gct gcc ttg aat
 1056
 Leu Val Gly Leu Ile Pro Ser Asn Ile Ser Ser Cys Ala Ala Leu Asn
 340 345 350
 45
 caa ttc aat gtt cat ggg aac ttc ttg agt gga gct gta cca ctt gaa
 1104
 Gln Phe Asn Val His Gly Asn Phe Leu Ser Gly Ala Val Pro Leu Glu
 355 360 365
 50
 ttc cgg aat ctt gga agc ttg act tat cta aat ctt tcc tca aac agt
 1152
 Phe Arg Asn Leu Gly Ser Leu Thr Tyr Leu Asn Leu Ser Ser Asn Ser
 370 375 380
 55
 ttc aag ggc aaa ata cct gct gag ctt ggc cat atc atc aat ctt gat
 1200

Phe Lys Gly Lys Ile Pro Ala Glu Leu Gly His Ile Ile Asn Leu Asp
 385 390 395 400
 5 aca ttg gat ctg tct ggc aac aat ttc tca ggc tca att cca tta aca
 1248
 Thr Leu Asp Leu Ser Gly Asn Asn Phe Ser Gly Ser Ile Pro Leu Thr
 405 410 415
 10 ctt ggt gat ctt gag cat ctt ctc atc tta aac ttg agc aga aat cat
 1296
 Leu Gly Asp Leu Glu His Leu Leu Ile Leu Asn Leu Ser Arg Asn His
 420 425 430
 15 ctg aat ggc aca ttg cct gca gaa ttc ggg aac ctc cga agc att cag
 1344
 Leu Asn Gly Thr Leu Pro Ala Glu Phe Gly Asn Leu Arg Ser Ile Gln
 435 440 445
 20 atc atc gat gtg tca ttt aat ttt ctt gcc ggt gtt att cca act gaa
 1392
 Ile Ile Asp Val Ser Phe Asn Phe Leu Ala Gly Val Ile Pro Thr Glu
 450 455 460
 25 ctt ggc cag ttg cag aac ata aac tct ctg ata ctg aac aac aac aag
 1440
 Leu Gly Gln Leu Gln Asn Ile Asn Ser Leu Ile Leu Asn Asn Asn Lys
 465 470 475 480
 30 att cat ggg aaa atc cct gat cag cta act aac tgc ttc agt ctt gcc
 1488
 Ile His Gly Lys Ile Pro Asp Gln Leu Thr Asn Cys Phe Ser Leu Ala
 485 490 495
 35 aat ctg aac atc tcc ttc aat aat ctt tct gga ata atc cca cct atg
 1536
 Asn Leu Asn Ile Ser Phe Asn Asn Leu Ser Gly Ile Ile Pro Pro Met
 500 505 510
 40 aag aac ttt aca cgt ttt tcc ccg gcc agc ttc ttt gga aat cca ttt
 1584
 Lys Asn Phe Thr Arg Phe Ser Pro Ala Ser Phe Phe Gly Asn Pro Phe
 515 520 525
 45 ctc tgc ggg aac tgg gtt gga tca atc tgt ggc cca tct tta cct aag
 1632
 Leu Cys Gly Asn Trp Val Gly Ser Ile Cys Gly Pro Ser Leu Pro Lys
 530 535 540
 50 tca caa gta ttc acc aga gtt gcc gtg att tgt atg gtt ctc ggt ttc
 1680
 Ser Gln Val Phe Thr Arg Val Ala Val Ile Cys Met Val Leu Gly Phe
 545 550 555 560
 55 atc act ctc ata tgc atg ata ttc att gcg gtt tac aag tca aag cag
 1728
 Ile Thr Leu Ile Cys Met Ile Phe Ile Ala Val Tyr Lys Ser Lys Gln
 565 570 575

cag aaa cca gtc ttg aaa ggc tct tca aaa caa cct gaa ggg tca acg
 1776
 5 Gln Lys Pro Val Leu Lys Gly Ser Ser Lys Gln Pro Glu Gly Ser Thr
 580 585 590

aag ctg gtg att ctt cac atg gac atg gct att cac acg ttt gat gat
 1824
 10 Lys Leu Val Ile Leu His Met Asp Met Ala Ile His Thr Phe Asp Asp
 595 600 605

atc atg aga gtt aca gaa aac ctc gat gag aaa tac atc att gga tac
 1872
 15 Ile Met Arg Val Thr Glu Asn Leu Asp Glu Lys Tyr Ile Ile Gly Tyr
 610 615 620

ggt gct tct agc aca gtt tac aag tgc acc tcc aaa act tcc cga cct
 1920
 20 Gly Ala Ser Ser Thr Val Tyr Lys Cys Thr Ser Lys Thr Ser Arg Pro
 625 630 635 640

att gcc att aag cga atc tac aat cag tat ccc agc aac ttc cgc gag
 1968
 25 Ile Ala Ile Lys Arg Ile Tyr Asn Gln Tyr Pro Ser Asn Phe Arg Glu
 645 650 655

ttt gaa aca gag ctc gag acc att ggg agc atc aga cac aga aac ata
 2016
 30 Phe Glu Thr Glu Leu Glu Thr Ile Gly Ser Ile Arg His Arg Asn Ile
 660 665 670

gta agc ttg cac gga tac gcc tta tct ccc ttt ggc aac ctc ctc ttc
 2064
 35 Val Ser Leu His Gly Tyr Ala Leu Ser Pro Phe Gly Asn Leu Leu Phe
 675 680 685

tac gac tac atg gaa aat ggc tct ctt tgg gat ctt ctc cat ggg cct
 2112
 40 Tyr Asp Tyr Met Glu Asn Gly Ser Leu Trp Asp Leu Leu His Gly Pro
 690 695 700

ggg aag aag gtg aag ctt gac tgg gaa aca agg ctg aag ata gct gtt
 2160
 45 Gly Lys Lys Val Lys Leu Asp Trp Glu Thr Arg Leu Lys Ile Ala Val
 705 710 715 720

gga gct gcg caa gga ctt gca tat ctt cac cat gac tgc aca cct agg
 2208
 50 Gly Ala Ala Gln Gly Leu Ala Tyr Leu His His Asp Cys Thr Pro Arg
 725 730 735

ata atc cat cga gac atc aag tca tca aac ata ctc ctt gat ggg aat
 2256
 55 Ile Ile His Arg Asp Ile Lys Ser Ser Asn Ile Leu' Leu Asp Gly Asn
 740 745 750

ttc gaa gcg cgt ttg tca gat ttt ggg att gcc aag agc ata cca gcc
 2304
 Phe Glu Ala Arg Leu Ser Asp Phe Gly Ile Ala Lys Ser Ile Pro Ala
 755 760 765
 5
 acc aaa act tat gct tca acc tat gtt ctt gga acc att gga tat att
 2352
 Thr Lys Thr Tyr Ala Ser Thr Tyr Val Leu Gly Thr Ile Gly Tyr Ile
 770 775 780
 10
 gac cca gag tat gct cga act tcg cgt ctg aac gag aag tct gat atc
 2400
 Asp Pro Glu Tyr Ala Arg Thr Ser Arg Leu Asn Glu Lys Ser Asp Ile
 785 790 795 800
 15
 tac agt ttc ggt att gtc ctt ctt gag ctt cta acc ggc aag aag gct
 2448
 Tyr Ser Phe Gly Ile Val Leu Leu Glu Leu Leu Thr Gly Lys Lys Ala
 805 810 815
 20
 gtg gat aac gag gcc aac ttg cat caa atg att cta tca aag gcg gat
 2496
 Val Asp Asn Glu Ala Asn Leu His Gln Met Ile Leu Ser Lys Ala Asp
 820 825 830
 25
 gat aac aca gta atg gaa gct gtt gat gca gag gtc tca gtg act tgc
 2544
 Asp Asn Thr Val Met Glu Ala Val Asp Ala Glu Val Ser Val Thr Cys
 835 840 845
 30
 atg gac tca gga cac atc aag aaa aca ttt cag cta gct ctc ttg tgc
 2592
 Met Asp Ser Gly His Ile Lys Lys Thr Phe Gln Leu Ala Leu Leu Cys
 850 855 860
 35
 acc aag cga aat cct ttg gag aga ccc acc atg cag gag gtc tct agg
 2640
 Thr Lys Arg Asn Pro Leu Glu Arg Pro Thr Met Gln Glu Val Ser Arg
 865 870 875 880
 40
 gtt ctg ctc tca ctt gtc ccg tct cca cct cca aag aag tta ccg tcg
 2688
 Val Leu Leu Ser Leu Val Pro Ser Pro Pro Pro Lys Lys Leu Pro Ser
 885 890 895
 45
 cct gca aaa gta cag gaa ggg gaa gaa cgg cgt gag agc cac tct tca
 2736
 Pro Ala Lys Val Gln Glu Gly Glu Glu Arg Arg Glu Ser His Ser Ser
 900 905 910
 50
 gat aca aca acc cca cag tgg ttt gtt cag ttc cgt gaa gat atc tcc
 2784
 Asp Thr Thr Thr Pro Gln Trp Phe Val Gln Phe Arg Glu Asp Ile Ser
 915 920 925
 55
 aaa agt agc tta taa
 2799

Lys Ser Ser Leu
930

5 <210> 8

<211> 932

<212> PRT

10

<213> Arabdopsis thaliana ERECTA homolog

15 <400> 8

Met Ala Ile Lys Ala Ser Phe Ser Asn Val Ala Asn Met Leu Leu Asp
1 5 10 15

20

Trp Asp Asp Val His Asn His Asp Phe Cys Ser Trp Arg Gly Val Phe
20 25 30

25

Cys Asp Asn Val Ser Leu Asn Val Val Ser Leu Asn Leu Ser Asn Leu
35 40 45

30

Asn Leu Gly Gly Glu Ile Ser Ser Ala Leu Gly Asp Leu Met Asn Leu
50 55 60

35

Gln Ser Ile Asp Leu Gln Gly Asn Lys Leu Gly Gly Gln Ile Pro Asp
65 70 75 80

Glu Ile Gly Asn Cys Val Ser Leu Ala Tyr Val Asp Phe Ser Thr Asn
85 90 95

40

Leu Leu Phe Gly Asp Ile Pro Phe Ser Ile Ser Lys Leu Lys Gln Leu
100 105 110

45

Glu Phe Leu Asn Leu Lys Asn Asn Gln Leu Thr Gly Pro Ile Pro Ala
115 120 125

50

Thr Leu Thr Gln Ile Pro Asn Leu Lys Thr Leu Asp Leu Ala Arg Asn
130 135 140

55

Gln Leu Thr Gly Glu Ile Pro Arg Leu Leu Tyr Trp Asn Glu Val Leu
145 150 155 160

Gln Tyr Leu Gly Leu Arg Gly Asn Met Leu Thr Gly Thr Leu Ser Pro

	165		170		175														
5	Asp	Met	Cys	Gln	Leu	Thr	Gly	Leu	Trp	Tyr	Phe	Asp	Val	Arg	Gly	Asn			
				180					185					190					
10	Asn	Leu	Thr	Gly	Thr	Ile	Pro	Glu	Ser	Ile	Gly	Asn	Cys	Thr	Ser	Phe			
				195				200					205						
15	Glu	Ile	Leu	Asp	Val	Ser	Tyr	Asn	Gln	Ile	Thr	Gly	Val	Ile	Pro	Tyr			
		210					215					220							
20	Asn	Ile	Gly	Phe	Leu	Gln	Val	Ala	Thr	Leu	Ser	Leu	Gln	Gly	Asn	Lys			
	225					230					235					240			
25	Leu	Thr	Gly	Arg	Ile	Pro	Glu	Val	Ile	Gly	Leu	Met	Gln	Ala	Leu	Ala			
					245					250					255				
30	Val	Leu	Asp	Leu	Ser	Asp	Asn	Glu	Leu	Thr	Gly	Pro	Ile	Pro	Pro	Ile			
				260					265					270					
35	Leu	Gly	Asn	Leu	Ser	Phe	Thr	Gly	Lys	Leu	Tyr	Leu	His	Gly	Asn	Lys			
			275					280					285						
40	Leu	Thr	Gly	Gln	Ile	Pro	Pro	Glu	Leu	Gly	Asn	Met	Ser	Arg	Leu	Ser			
		290				295						300							
45	Tyr	Leu	Gln	Leu	Asn	Asp	Asn	Glu	Leu	Val	Gly	Lys	Ile	Pro	Pro	Glu			
	305					310					315					320			
50	Leu	Gly	Lys	Leu	Glu	Gln	Leu	Phe	Glu	Leu	Asn	Leu	Ala	Asn	Asn	Asn			
				325					330					335					
55	Leu	Val	Gly	Leu	Ile	Pro	Ser	Asn	Ile	Ser	Ser	Cys	Ala	Ala	Leu	Asn			
				340					345					350					
60	Gln	Phe	Asn	Val	His	Gly	Asn	Phe	Leu	Ser	Gly	Ala	Val	Pro	Leu	Glu			
			355					360					365						
65	Phe	Arg	Asn	Leu	Gly	Ser	Leu	Thr	Tyr	Leu	Asn	Leu	Ser	Ser	Asn	Ser			
		370					375					380							
70	Phe	Lys	Gly	Lys	Ile	Pro	Ala	Glu	Leu	Gly	His	Ile	Ile	Asn	Leu	Asp			
	385					390					395					400			

Thr Leu Asp Leu Ser Gly Asn Asn Phe Ser Gly Ser Ile Pro Leu Thr
 405 410 415
 5

Leu Gly Asp Leu Glu His Leu Leu Ile Leu Asn Leu Ser Arg Asn His
 420 425 430

10

Leu Asn Gly Thr Leu Pro Ala Glu Phe Gly Asn Leu Arg Ser Ile Gln
 435 440 445

15

Ile Ile Asp Val Ser Phe Asn Phe Leu Ala Gly Val Ile Pro Thr Glu
 450 455 460

20

Leu Gly Gln Leu Gln Asn Ile Asn Ser Leu Ile Leu Asn Asn Asn Lys
 465 470 475 480

25

Ile His Gly Lys Ile Pro Asp Gln Leu Thr Asn Cys Phe Ser Leu Ala
 485 490 495

Asn Leu Asn Ile Ser Phe Asn Asn Leu Ser Gly Ile Ile Pro Pro Met
 500 505 510

30

Lys Asn Phe Thr Arg Phe Ser Pro Ala Ser Phe Phe Gly Asn Pro Phe
 515 520 525

35

Leu Cys Gly Asn Trp Val Gly Ser Ile Cys Gly Pro Ser Leu Pro Lys
 530 535 540

40

Ser Gln Val Phe Thr Arg Val Ala Val Ile Cys Met Val Leu Gly Phe
 545 550 555 560

45

Ile Thr Leu Ile Cys Met Ile Phe Ile Ala Val Tyr Lys Ser Lys Gln
 565 570 575

Gln Lys Pro Val Leu Lys Gly Ser Ser Lys Gln Pro Glu Gly Ser Thr
 580 585 590

50

Lys Leu Val Ile Leu His Met Asp Met Ala Ile His Thr Phe Asp Asp
 595 600 605

55

Ile Met Arg Val Thr Glu Asn Leu Asp Glu Lys Tyr Ile Ile Gly Tyr
 610 615 620

	Gly	Ala	Ser	Ser	Thr	Val	Tyr	Lys	Cys	Thr	Ser	Lys	Thr	Ser	Arg	Pro	625	630	635	640
5	Ile	Ala	Ile	Lys	Arg	Ile	Tyr	Asn	Gln	Tyr	Pro	Ser	Asn	Phe	Arg	Glu	645	650	655	
10	Phe	Glu	Thr	Glu	Leu	Glu	Thr	Ile	Gly	Ser	Ile	Arg	His	Arg	Asn	Ile	660	665	670	
15	Val	Ser	Leu	His	Gly	Tyr	Ala	Leu	Ser	Pro	Phe	Gly	Asn	Leu	Leu	Phe	675	680	685	
20	Tyr	Asp	Tyr	Met	Glu	Asn	Gly	Ser	Leu	Trp	Asp	Leu	Leu	His	Gly	Pro	690	695	700	
	Gly	Lys	Lys	Val	Lys	Leu	Asp	Trp	Glu	Thr	Arg	Leu	Lys	Ile	Ala	Val	705	710	715	720
25	Gly	Ala	Ala	Gln	Gly	Leu	Ala	Tyr	Leu	His	His	Asp	Cys	Thr	Pro	Arg	725	730	735	
30	Ile	Ile	His	Arg	Asp	Ile	Lys	Ser	Ser	Asn	Ile	Leu	Leu	Asp	Gly	Asn	740	745	750	
35	Phe	Glu	Ala	Arg	Leu	Ser	Asp	Phe	Gly	Ile	Ala	Lys	Ser	Ile	Pro	Ala	755	760	765	
40	Thr	Lys	Thr	Tyr	Ala	Ser	Thr	Tyr	Val	Leu	Gly	Thr	Ile	Gly	Tyr	Ile	770	775	780	
	Asp	Pro	Glu	Tyr	Ala	Arg	Thr	Ser	Arg	Leu	Asn	Glu	Lys	Ser	Asp	Ile	785	790	795	800
45	Tyr	Ser	Phe	Gly	Ile	Val	Leu	Leu	Glu	Leu	Leu	Thr	Gly	Lys	Lys	Ala	805	810	815	
50	Val	Asp	Asn	Glu	Ala	Asn	Leu	His	Gln	Met	Ile	Leu	Ser	Lys	Ala	Asp	820	825	830	
55	Asp	Asn	Thr	Val	Met	Glu	Ala	Val	Asp	Ala	Glu	Val	Ser	Val	Thr	Cys	835	840	845	

Met Asp Ser Gly His Ile Lys Lys Thr Phe Gln Leu Ala Leu Leu Cys
850 855 860

5 Thr Lys Arg Asn Pro Leu Glu Arg Pro Thr Met Gln Glu Val Ser Arg
865 870 875 880

10 Val Leu Leu Ser Leu Val Pro Ser Pro Pro Pro Lys Lys Leu Pro Ser
885 890 895

15 Pro Ala Lys Val Gln Glu Gly Glu Glu Arg Arg Glu Ser His Ser Ser
900 905 910

20 Asp Thr Thr Thr Pro Gln Trp Phe Val Gln Phe Arg Glu Asp Ile Ser
915 920 925

Lys Ser Ser Leu
930

25 <210> 9
<211> 2901
<212> DNA
30 <213> Aradopsis thaliana ERECTA homolog

35 <220>
<221> CDS
<222> (1)..(2898)
40 <223>

45 <400> 9
atg aag gag aag atg cag cga atg gtt tta tct tta gca atg gtg ggt
48
Met Lys Glu Lys Met Gln Arg Met Val Leu Ser Leu Ala Met Val Gly
1 5 10 15
50 ttt atg gtt ttt ggt gtt gct tcg gct atg aac aac gaa ggg aaa gct
96
Phe Met Val Phe Gly Val Ala Ser Ala Met Asn Asn Glu Gly Lys Ala
20 25 30
55 ctg atg gcg ata aaa ggc tct ttc agc aac tta gtg aat atg ctt ttg
144

	Leu	Met	Ala	Ile	Lys	Gly	Ser	Phe	Ser	Asn	Leu	Val	Asn	Met	Leu	Leu
			35					40					45			
5	gat	tgg	gac	gat	gtt	cac	aac	agt	gac	ttg	tgt	tct	tgg	cga	ggt	gtt
	192															
	Asp	Trp	Asp	Asp	Val	His	Asn	Ser	Asp	Leu	Cys	Ser	Trp	Arg	Gly	Val
		50					55					60				
10	ttc	tgc	gac	aac	gtt	agc	tac	tcc	gtt	gtc	tct	ctg	aat	ttg	tcc	agt
	240															
	Phe	Cys	Asp	Asn	Val	Ser	Tyr	Ser	Val	Val	Ser	Leu	Asn	Leu	Ser	Ser
	65					70					75					80
15	ctg	aat	ctt	gga	ggg	gag	ata	tct	cca	gct	att	gga	gac	cta	cgg	aat
	288															
	Leu	Asn	Leu	Gly	Gly	Glu	Ile	Ser	Pro	Ala	Ile	Gly	Asp	Leu	Arg	Asn
					85					90						95
20	ttg	caa	tca	ata	gac	ttg	caa	ggt	aat	aaa	cta	gca	ggt	caa	att	cca
	336															
	Leu	Gln	Ser	Ile	Asp	Leu	Gln	Gly	Asn	Lys	Leu	Ala	Gly	Gln	Ile	Pro
				100					105						110	
25	gat	gag	att	gga	aac	tgt	gct	tct	ctt	gtt	tat	ctg	gat	ttg	tcc	gag
	384															
	Asp	Glu	Ile	Gly	Asn	Cys	Ala	Ser	Leu	Val	Tyr	Leu	Asp	Leu	Ser	Glu
			115					120					125			
30	aat	ctg	tta	tat	gga	gac	ata	cct	ttc	tca	atc	tct	aaa	ctc	aag	cag
	432															
	Asn	Leu	Leu	Tyr	Gly	Asp	Ile	Pro	Phe	Ser	Ile	Ser	Lys	Leu	Lys	Gln
		130					135						140			
35	ctt	gaa	act	ctg	aat	ctg	aag	aac	aat	cag	ctc	aca	ggt	cct	gta	cca
	480															
	Leu	Glu	Thr	Leu	Asn	Leu	Lys	Asn	Asn	Gln	Leu	Thr	Gly	Pro	Val	Pro
	145					150					155					160
40	gca	acc	tta	acc	cag	att	cca	aac	ctt	aag	aga	ctt	gat	ctt	gct	ggc
	528															
	Ala	Thr	Leu	Thr	Gln	Ile	Pro	Asn	Leu	Lys	Arg	Leu	Asp	Leu	Ala	Gly
					165					170					175	
45	aat	cat	cta	acg	ggt	gag	ata	tcg	aga	ttg	ctt	tac	tgg	aat	gaa	gtt
	576															
	Asn	His	Leu	Thr	Gly	Glu	Ile	Ser	Arg	Leu	Leu	Tyr	Trp	Asn	Glu	Val
				180					185					190		
50	ttg	cag	tat	ctt	gga	tta	cga	ggg	aat	atg	ttg	act	gga	acg	tta	tct
	624															
	Leu	Gln	Tyr	Leu	Gly	Leu	Arg	Gly	Asn	Met	Leu	Thr	Gly	Thr	Leu	Ser
			195					200					205			
55	tct	gat	atg	tgt	cag	cta	acc	ggt	ttg	tgg	tac	ttt	gat	gtg	aga	gga
	672															
	Ser	Asp	Met	Cys	Gln	Leu	Thr	Gly	Leu	Trp	Tyr	Phe	Asp	Val	Arg	Gly
		210					215					220				

	aat aat cta act gga acc atc ccg gag agc atc gga aat tgc aca agc	
	720	
5	Asn Asn Leu Thr Gly Thr Ile Pro Glu Ser Ile Gly Asn Cys Thr Ser	
	225 230 235 240	
	ttt caa atc ctg gac ata tct tat aat cag ata aca gga gag att cct	
	768	
10	Phe Gln Ile Leu Asp Ile Ser Tyr Asn Gln Ile Thr Gly Glu Ile Pro	
	245 250 255	
	tac aat atc ggc ttc ctc caa gtt gct act ctg tca ctt caa gga aac	
	816	
15	Tyr Asn Ile Gly Phe Leu Gln Val Ala Thr Leu Ser Leu Gln Gly Asn	
	260 265 270	
	aga ttg acg ggt aga att cca gaa gtt att ggt cta atg cag gct ctt	
	864	
20	Arg Leu Thr Gly Arg Ile Pro Glu Val Ile Gly Leu Met Gln Ala Leu	
	275 280 285	
	gct gtt ttg gat ttg agt gac aat gag ctt gtt ggt cct atc cca ccg	
	912	
25	Ala Val Leu Asp Leu Ser Asp Asn Glu Leu Val Gly Pro Ile Pro Pro	
	290 295 300	
	ata ctt ggc aat ctc tca ttt acc gga aag ttg tat ctc cat ggc aat	
	960	
30	Ile Leu Gly Asn Leu Ser Phe Thr Gly Lys Leu Tyr Leu His Gly Asn	
	305 310 315 320	
	atg ctc act ggt cca atc ccc tct gag ctt ggg aat atg tca cgt ctc	
	1008	
35	Met Leu Thr Gly Pro Ile Pro Ser Glu Leu Gly Asn Met Ser Arg Leu	
	325 330 335	
	agc tat ttg cag cta aac gac aat aaa cta gtg gga act att cca cct	
	1056	
40	Ser Tyr Leu Gln Leu Asn Asp Asn Lys Leu Val Gly Thr Ile Pro Pro	
	340 345 350	
	gag ctt gga aag ctg gag caa ttg ttt gaa ctg aat ctt gcc aac aac	
	1104	
45	Glu Leu Gly Lys Leu Glu Gln Leu Phe Glu Leu Asn Leu Ala Asn Asn	
	355 360 365	
	cgt tta gta ggg ccc ata cca tcc aac att agt tca tgt gca gcc ttg	
	1152	
50	Arg Leu Val Gly Pro Ile Pro Ser Asn Ile Ser Ser Cys Ala Ala Leu	
	370 375 380	
	aat caa ttc aat gtt cat ggg aac ctc ttg agt gga tct att cca ctg	
	1200	
55	Asn Gln Phe Asn Val His Gly Asn Leu Leu Ser Gly Ser Ile Pro Leu	
	385 390 395 400	

gcg ttt cgc aat ctc ggg agc ttg act tat ctg aat ctt tcg tcg aac
1248
Ala Phe Arg Asn Leu Gly Ser Leu Thr Tyr Leu Asn Leu Ser Ser Asn
405 410 415

5 aat ttc aag gga aaa ata cca gtt gag ctt gga cat ata atc aat ctt
1296
Asn Phe Lys Gly Lys Ile Pro Val Glu Leu Gly His Ile Ile Asn Leu
420 425 430

10 gac aaa cta gat ctg tct ggc aat aac ttc tca ggg tct ata cca tta
1344
Asp Lys Leu Asp Leu Ser Gly Asn Asn Phe Ser Gly Ser Ile Pro Leu
435 440 445

15 acg ctt ggc gat ctt gaa cac ctt ctc ata tta aat ctt agc aga aac
1392
Thr Leu Gly Asp Leu Glu His Leu Leu Ile Leu Asn Leu Ser Arg Asn
450 455 460

20 cat ctt agt gga caa tta cct gca gag ttt ggg aac ctt cga agc att
1440
His Leu Ser Gly Gln Leu Pro Ala Glu Phe Gly Asn Leu Arg Ser Ile
465 470 475 480

25 cag atg att gat gta tca ttc aat ctg ctc tcc gga gtt att cca act
1488
Gln Met Ile Asp Val Ser Phe Asn Leu Leu Ser Gly Val Ile Pro Thr
485 490 495

30 gaa ctt ggc caa ttg cag aat tta aac tct tta ata ttg aac aac aac
1536
Glu Leu Gly Gln Leu Gln Asn Leu Asn Ser Leu Ile Leu Asn Asn Asn
500 505 510

35 aag ctt cat ggg aaa att cca gat cag ctt acg aac tgc ttc act ctt
1584
Lys Leu His Gly Lys Ile Pro Asp Gln Leu Thr Asn Cys Phe Thr Leu
515 520 525

40 gtc aat ctg aat gtc tcc ttc aac aat ctc tcc ggg ata gtc cca cca
1632
Val Asn Leu Asn Val Ser Phe Asn Asn Leu Ser Gly Ile Val Pro Pro
530 535 540

45 atg aaa aac ttc tca cgt ttt gct cca gcc agc ttt gtt gga aat cca
1680
Met Lys Asn Phe Ser Arg Phe Ala Pro Ala Ser Phe Val Gly Asn Pro
545 550 555 560

50 tat ctt tgt gga aac tgg gtt gga tct att tgt ggt cct tta ccg aaa
1728
Tyr Leu Cys Gly Asn Trp Val Gly Ser Ile Cys Gly Pro Leu Pro Lys
565 570 575

55 tct cga gta ttc tcc aga ggt gct ttg atc tgc att gtt ctt ggc gtc
1776

	Ser	Arg	Val	Phe	Ser	Arg	Gly	Ala	Leu	Ile	Cys	Ile	Val	Leu	Gly	Val
				580					585					590		
5	atc	act	ctc	cta	tgt	atg	att	ttc	ctt	gca	ggt	tac	aaa	tca	atg	cag
	1824															
	Ile	Thr	Leu	Leu	Cys	Met	Ile	Phe	Leu	Ala	Val	Tyr	Lys	Ser	Met	Gln
			595					600					605			
10	cag	aag	aag	att	cta	caa	ggc	tcc	tca	aaa	caa	gct	gaa	ggg	tta	acc
	1872															
	Gln	Lys	Lys	Ile	Leu	Gln	Gly	Ser	Ser	Lys	Gln	Ala	Glu	Gly	Leu	Thr
		610					615					620				
15	aag	cta	gtg	att	ctc	cac	atg	gac	atg	gca	att	cat	aca	ttt	gat	gat
	1920															
	Lys	Leu	Val	Ile	Leu	His	Met	Asp	Met	Ala	Ile	His	Thr	Phe	Asp	Asp
	625					630					635				640	
20	atc	atg	aga	gtg	act	gag	aat	ctt	aac	gaa	aag	ttt	ata	att	gga	tat
	1968															
	Ile	Met	Arg	Val	Thr	Glu	Asn	Leu	Asn	Glu	Lys	Phe	Ile	Ile	Gly	Tyr
					645					650					655	
25	ggc	gct	tct	agc	acg	gta	tac	aaa	tgt	gca	tta	aaa	agt	tcc	cga	cct
	2016															
	Gly	Ala	Ser	Ser	Thr	Val	Tyr	Lys	Cys	Ala	Leu	Lys	Ser	Ser	Arg	Pro
				660					665					670		
30	att	gcc	att	aag	cga	ctc	tac	aat	cag	tat	ccg	cat	aac	ttg	cgg	gaa
	2064															
	Ile	Ala	Ile	Lys	Arg	Leu	Tyr	Asn	Gln	Tyr	Pro	His	Asn	Leu	Arg	Glu
			675					680					685			
35	ttt	gag	aca	gaa	ctt	gag	acc	att	ggg	agc	att	agg	cac	aga	aac	ata
	2112															
	Phe	Glu	Thr	Glu	Leu	Glu	Thr	Ile	Gly	Ser	Ile	Arg	His	Arg	Asn	Ile
		690					695					700				
40	gtc	agc	ttg	cat	gga	tat	gcc	ttg	tct	cct	act	ggc	aac	ctt	ctt	ttc
	2160															
	Val	Ser	Leu	His	Gly	Tyr	Ala	Leu	Ser	Pro	Thr	Gly	Asn	Leu	Leu	Phe
	705					710					715					720
45	tat	gac	tac	atg	gaa	aat	gga	tca	ctt	tgg	gac	ctt	ctt	cat	ggg	tca
	2208															
	Tyr	Asp	Tyr	Met	Glu	Asn	Gly	Ser	Leu	Trp	Asp	Leu	Leu	His	Gly	Ser
					725					730					735	
50	ttg	aag	aaa	gtg	aag	ctt	gat	tgg	gag	aca	agg	ttg	aag	ata	gcg	gtt
	2256															
	Leu	Lys	Lys	Val	Lys	Leu	Asp	Trp	Glu	Thr	Arg	Leu	Lys	Ile	Ala	Val
				740					745					750		
55	gga	gct	gca	caa	gga	cta	gcc	tat	ctt	cac	cac	gat	tgt	act	cct	cga
	2304															
	Gly	Ala	Ala	Gln	Gly	Leu	Ala	Tyr	Leu	His	His	Asp	Cys	Thr	Pro	Arg
			755					760					765			

atc att cac cgt gac atc aag tca tcg aac ata ctt ctt gat gag aat
 2352
 5 Ile Ile His Arg Asp Ile Lys Ser Ser Asn Ile Leu Leu Asp Glu Asn
 770 775 780

ttc gaa gca cat tta tct gat ttc ggg att gct aag agc ata cca gct
 2400
 10 Phe Glu Ala His Leu Ser Asp Phe Gly Ile Ala Lys Ser Ile Pro Ala
 785 790 795 800

agc aaa acc cat gcc tcg act tat gtt ttg gga aca att ggt tat ata
 2448
 15 Ser Lys Thr His Ala Ser Thr Tyr Val Leu Gly Thr Ile Gly Tyr Ile
 805 810 815

gac cca gag tat gct cgt act tca cga atc aat gag aaa tcc gat ata
 2496
 20 Asp Pro Glu Tyr Ala Arg Thr Ser Arg Ile Asn Glu Lys Ser Asp Ile
 820 825 830

tac agc ttc ggt att gtt ctt ctt gag ctt ctc act ggg aag aaa gca
 2544
 25 Tyr Ser Phe Gly Ile Val Leu Leu Glu Leu Leu Thr Gly Lys Lys Ala
 835 840 845

gtg gat aac gaa gct aac ttg cat caa ctg ata ttg tca aag gct gat
 2592
 30 Val Asp Asn Glu Ala Asn Leu His Gln Leu Ile Leu Ser Lys Ala Asp
 850 855 860

gat aat act gtg atg gaa gca gtt gat cca gag gtt act gtg act tgt
 2640
 35 Asp Asn Thr Val Met Glu Ala Val Asp Pro Glu Val Thr Val Thr Cys
 865 870 875 880

atg gac ttg gga cat atc agg aag aca ttt cag ctg gct ctc tta tgc
 2688
 40 Met Asp Leu Gly His Ile Arg Lys Thr Phe Gln Leu Ala Leu Leu Cys
 885 890 895

aca aag cga aac cct tta gag aga ccc aca atg ctt gaa gtc tct agg
 2736
 45 Thr Lys Arg Asn Pro Leu Glu Arg Pro Thr Met Leu Glu Val Ser Arg
 900 905 910

gtt ctg ctc tct ctt gtc cca tct ctg caa gta gca aag aag cta cct
 2784
 50 Val Leu Leu Ser Leu Val Pro Ser Leu Gln Val Ala Lys Lys Leu Pro
 915 920 925

tct ctt gat cac tca acc aaa aag ctg cag caa gag aat gaa gtt agg
 2832
 55 Ser Leu Asp His Ser Thr Lys Lys Leu Gln Gln Glu Asn Glu Val Arg
 930 935 940

aat cct gat gca gaa gca tct caa tgg ttt gtt cag ttc cgt gaa gtc
 2880
 Asn Pro Asp Ala Glu Ala Ser Gln Trp Phe Val Gln Phe Arg Glu Val
 945 950 955 960
 5
 atc tcc aaa agt agc ata taa
 2901
 Ile Ser Lys Ser Ser Ile
 965
 10
 <210> 10
 <211> 966
 15
 <212> PRT
 <213> Aradopsis thaliana ERECTA homolog
 20
 <400> 10
 25 Met Lys Glu Lys Met Gln Arg Met Val Leu Ser Leu Ala Met Val Gly
 1 5 10 15
 Phe Met Val Phe Gly Val Ala Ser Ala Met Asn Asn Glu Gly Lys Ala
 20 25 30
 30 Leu Met Ala Ile Lys Gly Ser Phe Ser Asn Leu Val Asn Met Leu Leu
 35 40 45
 35 Asp Trp Asp Asp Val His Asn Ser Asp Leu Cys Ser Trp Arg Gly Val
 50 55 60
 40 Phe Cys Asp Asn Val Ser Tyr Ser Val Val Ser Leu Asn Leu Ser Ser
 65 70 75 80
 45 Leu Asn Leu Gly Gly Glu Ile Ser Pro Ala Ile Gly Asp Leu Arg Asn
 85 90 95
 Leu Gln Ser Ile Asp Leu Gln Gly Asn Lys Leu Ala Gly Gln Ile Pro
 100 105 110
 50 Asp Glu Ile Gly Asn Cys Ala Ser Leu Val Tyr Leu Asp Leu Ser Glu
 115 120 125
 55 Asn Leu Leu Tyr Gly Asp Ile Pro Phe Ser Ile Ser Lys Leu Lys Gln
 130 135 140

5 Leu Glu Thr Leu Asn Leu Lys Asn Asn Gln Leu Thr Gly Pro Val Pro
 145 150 155 160

10 Ala Thr Leu Thr Gln Ile Pro Asn Leu Lys Arg Leu Asp Leu Ala Gly
 165 170 175

15 Asn His Leu Thr Gly Glu Ile Ser Arg Leu Leu Tyr Trp Asn Glu Val
 180 185 190

20 Leu Gln Tyr Leu Gly Leu Arg Gly Asn Met Leu Thr Gly Thr Leu Ser
 195 200 205

25 Ser Asp Met Cys Gln Leu Thr Gly Leu Trp Tyr Phe Asp Val Arg Gly
 210 215 220

30 Asn Asn Leu Thr Gly Thr Ile Pro Glu Ser Ile Gly Asn Cys Thr Ser
 225 230 235 240

35 Phe Gln Ile Leu Asp Ile Ser Tyr Asn Gln Ile Thr Gly Glu Ile Pro
 245 250 255

40 Tyr Asn Ile Gly Phe Leu Gln Val Ala Thr Leu Ser Leu Gln Gly Asn
 260 265 270

45 Arg Leu Thr Gly Arg Ile Pro Glu Val Ile Gly Leu Met Gln Ala Leu
 275 280 285

50 Ala Val Leu Asp Leu Ser Asp Asn Glu Leu Val Gly Pro Ile Pro Pro
 290 295 300

55 Ile Leu Gly Asn Leu Ser Phe Thr Gly Lys Leu Tyr Leu His Gly Asn
 305 310 315 320

60 Met Leu Thr Gly Pro Ile Pro Ser Glu Leu Gly Asn Met Ser Arg Leu
 325 330 335

65 Ser Tyr Leu Gln Leu Asn Asp Asn Lys Leu Val Gly Thr Ile Pro Pro
 340 345 350

70 Glu Leu Gly Lys Leu Glu Gln Leu Phe Glu Leu Asn Leu Ala Asn Asn
 355 360 365

	Arg	Leu	Val	Gly	Pro	Ile	Pro	Ser	Asn	Ile	Ser	Ser	Cys	Ala	Ala	Leu	
	370						375					380					
5	Asn	Gln	Phe	Asn	Val	His	Gly	Asn	Leu	Leu	Ser	Gly	Ser	Ile	Pro	Leu	
	385					390					395					400	
10	Ala	Phe	Arg	Asn	Leu	Gly	Ser	Leu	Thr	Tyr	Leu	Asn	Leu	Ser	Ser	Asn	
					405					410						415	
15	Asn	Phe	Lys	Gly	Lys	Ile	Pro	Val	Glu	Leu	Gly	His	Ile	Ile	Asn	Leu	
				420					425							430	
20	Asp	Lys	Leu	Asp	Leu	Ser	Gly	Asn	Asn	Phe	Ser	Gly	Ser	Ile	Pro	Leu	
			435					440						445			
	Thr	Leu	Gly	Asp	Leu	Glu	His	Leu	Leu	Ile	Leu	Asn	Leu	Ser	Arg	Asn	
	450						455						460				
25	His	Leu	Ser	Gly	Gln	Leu	Pro	Ala	Glu	Phe	Gly	Asn	Leu	Arg	Ser	Ile	
	465					470					475					480	
30	Gln	Met	Ile	Asp	Val	Ser	Phe	Asn	Leu	Leu	Ser	Gly	Val	Ile	Pro	Thr	
					485					490						495	
35	Glu	Leu	Gly	Gln	Leu	Gln	Asn	Leu	Asn	Ser	Leu	Ile	Leu	Asn	Asn	Asn	
				500					505							510	
40	Lys	Leu	His	Gly	Lys	Ile	Pro	Asp	Gln	Leu	Thr	Asn	Cys	Phe	Thr	Leu	
			515					520						525			
45	Val	Asn	Leu	Asn	Val	Ser	Phe	Asn	Asn	Leu	Ser	Gly	Ile	Val	Pro	Pro	
		530					535						540				
	Met	Lys	Asn	Phe	Ser	Arg	Phe	Ala	Pro	Ala	Ser	Phe	Val	Gly	Asn	Pro	
	545					550					555					560	
50	Tyr	Leu	Cys	Gly	Asn	Trp	Val	Gly	Ser	Ile	Cys	Gly	Pro	Leu	Pro	Lys	
					565					570						575	
55	Ser	Arg	Val	Phe	Ser	Arg	Gly	Ala	Leu	Ile	Cys	Ile	Val	Leu	Gly	Val	
				580					585							590	

	Ile	Thr	Leu	Leu	Cys	Met	Ile	Phe	Leu	Ala	Val	Tyr	Lys	Ser	Met	Gln
			595					600					605			
5	Gln	Lys	Lys	Ile	Leu	Gln	Gly	Ser	Ser	Lys	Gln	Ala	Glu	Gly	Leu	Thr
	610						615					620				
10	Lys	Leu	Val	Ile	Leu	His	Met	Asp	Met	Ala	Ile	His	Thr	Phe	Asp	Asp
	625					630					635					640
15	Ile	Met	Arg	Val	Thr	Glu	Asn	Leu	Asn	Glu	Lys	Phe	Ile	Ile	Gly	Tyr
					645					650					655	
20	Gly	Ala	Ser	Ser	Thr	Val	Tyr	Lys	Cys	Ala	Leu	Lys	Ser	Ser	Arg	Pro
				660					665						670	
25	Ile	Ala	Ile	Lys	Arg	Leu	Tyr	Asn	Gln	Tyr	Pro	His	Asn	Leu	Arg	Glu
		675						680					685			
30	Phe	Glu	Thr	Glu	Leu	Glu	Thr	Ile	Gly	Ser	Ile	Arg	His	Arg	Asn	Ile
	690						695					700				
35	Val	Ser	Leu	His	Gly	Tyr	Ala	Leu	Ser	Pro	Thr	Gly	Asn	Leu	Leu	Phe
	705				710						715					720
40	Tyr	Asp	Tyr	Met	Glu	Asn	Gly	Ser	Leu	Trp	Asp	Leu	Leu	His	Gly	Ser
					725					730					735	
45	Leu	Lys	Lys	Val	Lys	Leu	Asp	Trp	Glu	Thr	Arg	Leu	Lys	Ile	Ala	Val
				740					745					750		
50	Gly	Ala	Ala	Gln	Gly	Leu	Ala	Tyr	Leu	His	His	Asp	Cys	Thr	Pro	Arg
		755					760						765			
55	Ile	Ile	His	Arg	Asp	Ile	Lys	Ser	Ser	Asn	Ile	Leu	Leu	Asp	Glu	Asn
	770					775						780				
60	Phe	Glu	Ala	His	Leu	Ser	Asp	Phe	Gly	Ile	Ala	Lys	Ser	Ile	Pro	Ala
	785					790					795					800
65	Ser	Lys	Thr	His	Ala	Ser	Thr	Tyr	Val	Leu	Gly	Thr	Ile	Gly	Tyr	Ile
				805						810				815		
70	Asp	Pro	Glu	Tyr	Ala	Arg	Thr	Ser	Arg	Ile	Asn	Glu	Lys	Ser	Asp	Ile

	820		825		830												
5	Tyr	Ser	Phe	Gly	Ile	Val	Leu	Leu	Glu	Leu	Leu	Thr	Gly	Lys	Lys	Ala	
			835					840					845				
10	Val	Asp	Asn	Glu	Ala	Asn	Leu	His	Gln	Leu	Ile	Leu	Ser	Lys	Ala	Asp	
		850					855					860					
15	Asp	Asn	Thr	Val	Met	Glu	Ala	Val	Asp	Pro	Glu	Val	Thr	Val	Thr	Cys	
	865					870					875					880	
20	Met	Asp	Leu	Gly	His	Ile	Arg	Lys	Thr	Phe	Gln	Leu	Ala	Leu	Leu	Cys	
					885					890					895		
25	Thr	Lys	Arg	Asn	Pro	Leu	Glu	Arg	Pro	Thr	Met	Leu	Glu	Val	Ser	Arg	
				900					905					910			
30	Val	Leu	Leu	Ser	Leu	Val	Pro	Ser	Leu	Gln	Val	Ala	Lys	Lys	Leu	Pro	
			915					920					925				
35	Ser	Leu	Asp	His	Ser	Thr	Lys	Lys	Leu	Gln	Gln	Glu	Asn	Glu	Val	Arg	
	930						935					940					
40	Asn	Pro	Asp	Ala	Glu	Ala	Ser	Gln	Trp	Phe	Val	Gln	Phe	Arg	Glu	Val	
	945					950					955					960	
	Ile	Ser	Lys	Ser	Ser	Ile											
					965												

Dated this SECOND day of JULY, 2002

The Australian National University
Patent Attorneys for the Applicant:

F B RICE & CO




Figure 1a




Figure 1b




Figure 2a




Figure 2b




Figure 1c




Figure 2c